

The Relation of Alcohol Consumption and Blood Cholesterol Level

Joice Nancy Ansye Engka*, Hedison Polii, Sylvia Ritta Marunduh

Faculty of Medicine, Sam Ratulangi University Jl Kampus Unsrat, Bahu Manado 95115, Indonesia

Original Research Article

*Corresponding author

Joice Nancy Ansye Engka

Article History

Received: 27.11.2018

Accepted: 06.12.2018

Published: 30.12.2018

DOI:

10.36347/sjams.2018.v06i12.001



Abstract: Coronary Heart Disease is usually caused by atherosclerosis. Atherosclerosis can be developed because of the increase of blood cholesterol level. According to *World Health Organization* (WHO) in 2012 heart and vascular disease caused 17.5 million equal to 31% death around the world and 7.4 million was caused by Coronary Heart Disease. Center of health research/Riset kesehatan dasar (Riskesmas) shows in 2007 there are largest numbers of male alcohol drinkers aged 15 years old and older; 31.1% in North Sulawesi suburban area and 32.9% in countryside. General objective is to determine the relation between alcohol intake with blood cholesterol level with specific aim to study the relation between duration and volume of daily alcohol intake with cholesterol level. This research is observational analysis with cross sectional study. The subjects are 51 male students of Faculty of Engineering of Sam Ratulangi University aged 19 – 22 year old. Normal cholesterol level is 200 mg/dL. The result is there is a significant association between the duration of alcohol intake with cholesterol level ($p < 0.05$) and there is also a significant relation between the average daily intake ($p < 0.05$) with cholesterol level. Alcohol intake can influence blood cholesterol level.

Keywords: Alcohol intake, blood cholesterol.

INTRODUCTION

Coronary Heart Disease (CHD) is usually caused by atherosclerosis. Atherosclerosis can be caused by the increase of cholesterol level and results in pathologic accumulation in blood vessels [1]. High level of total blood cholesterol causes increasing in morbidity and mortality rates of cardiovascular disorders.

Based on prevalence men with high cholesterol level have higher risk to develop cardiovascular diseases. According to *World Health Organization* (WHO), in 2012 there are 17.5 million equal to 31 % deaths worldwide caused by Cardiovascular diseases and 7.4 millions of it caused by CHD. Based on doctor's diagnosis the prevalence of CHD in Indonesia in 2013 on the age of ≥ 15 year old is 0, 5% equal with 883.447 people, while based on symptoms is 1, 5% or about 2.650.340 people. Based on doctors diagnosis the largest number of people with CHD is in West Jawa province 160.812 orang (0, 5%). North Sulawesi has 11.892 cases (0.7%) based on doctor diagnosis and 28.880 cases (1, 7%) based on symptoms [3].

Nowadays alcohol consumption is becoming trending issue in several areas in Indonesia. Center of health research/Riset kesehatan dasar (Riskesmas) shows in 2007 there are largest numbers of male alcohol drinkers aged 15 years old and older; 31.1% in

North Sulawesi suburban area and 32.9% in countryside [4].

In the global era when lifestyle and eating pattern are changing, the level of blood cholesterol becomes higher in productive aged people as well as in late teenage aged 17-25 tahun [5]. Therefore it is necessary for the author to determine the effect of alcohol intake on blood cholesterol levels on university students.

MATERIALS AND METHODS

This research is a 10 months observational study with a cross sectional design. The subjects are 51 male students aged 19 – 22 years old of Faculty of Engineering Sam Ratulangi University badge 2015 and 2016 who already signed the informed consent. Blood were obtained from each subject and analysed for cholesterol levels with normal reference of 200 mg/dL. After that the association were determined by statistic analysis.

RESULTS AND DISCUSSION

Table-1: The correlation between alcohol consumption and cholesterol level

Duration alcohol consumption	Cholesterol Level				p
	Normal		Hypercholesterol		
	n	%	n	%	
< 3 year	29	74.4	8	66.7	0,023
≥ 3 year	10	25.6	4	33.3	
Total	39	100	12	100	

Based on table 1 there is a connection between alcohol intake duration and cholesterol levels ($p < 0,05$).

Table 2 shows a connection between the average daily intake and cholesterol levels ($p < 0.05$).

The subjects on this study are university students year 3 and year 4 aged 19 – 22 years old which

categorized as late teenage [4]. This certain age environmental factor is strongly influence the behavior of alcohol consumption. A study of Wijaya I.P.A on male teenagers in Bali stated there is a significant relation between alcohol intake and the social behavior between those teenagers [6]. There is also a similar study by Adiputra and Saputro with the similar result [7, 8].

Table-2: The correlation between the average daily intake (volume in mL) and cholesterol level

The average daily intake (volume in mL)	Cholesterol Level				P
	Normal		Hypercholesterol		
	n	%	n	%	
< 90 mL	18	46.2	0	35.3	0,010
≥ 90 mL	21	53.8	12	64.7	
Total	39	100	12	100	

Research by De Oliveira e Silva ER *et al* in 2000 stated that alcohol consumption increases cholesterol levels [9] also a study from Wakabayashi I, in 2010 and Yoon Y S *et al.* in 2004 found a significant relationship between alcohol consumption and cholesterol levels[10,11]. Research by Huang *et al.* in 2017 moderate alcohol consumption lowers blood cholesterol [12] as well as research conducted by Vu. Khanh *et al.* in 2016 alcohol consumption low to moderate can significantly increase HDL and reduce triglycerides, total cholesterol and LDL [13].

CONCLUSION

Alcohol consumption can influence blood cholesterol levels.

ACKNOWLEDGEMENT

A special thank you to Sam Ratulangi University, to the Dean and Vice Dean and the students badge 2015/2016 of Faculty of Engineering.

REFERENCES

1. American Heart Association.. Prevention and Treatment Options for Dislipidemi. 2008.
2. World Health Organization.Global Status Report on Alcohol and Health.WHO press.2011
3. Health Research and Development Agency of the Ministry of Health of the Republic of Indonesia. Research Report on Basic Health (Riskesdas) Indonesia. Jakarta. 2013.

4. Health Research and Development Agency of the Ministry of Health of the Republic of Indonesia Research Report on Basic Health (Riskesdas) Indonesia. Jakarta. 2007.
5. Yusuf, Syamsu. *Psikologi Perkembangan Anak dan Remaja*. Bandung: PT Remaja Rusdakarya. 2007.
6. Wijaya A. Faktor-faktor Yang Mempengaruhi Tingginya Konsumsi Alkohol Pada Remaja Putra Di Desa Keramas Kecamatan Blahbatuh Kabupaten Gianyar. *Jurnal Dunia Kesehatan*. 2016;5(2).
7. I Made Sudarma Adiputra, Siluh Nyoman Alit Nuryani, Dewa Ayu PutriAnggi Wirani.2008.Anlysis of Factors That Associated With an Incidence of Alcohol Consumption. *STIKES Wira Medika PPNI Bali, RSUP Sanglah Denpasar Prov. Bali,Jurnal Keperawatan FIKKeS*.2008
8. Fadli Alam Dwi Saputro,Yuni Dwi Hastuti,Triana Arisdiani, *Pengaruh Teman Sebaya Terhadap Prilaku Konsumsi Alkohol Pada Remaja Putra*. *JurnalIlmia STIKES Kendal*. 2014; 4(2)70-81.
9. De Oliveira e Silva ER, Foster D, McGee Harper M, Seidman CE, Smith JD, Breslow JL, Brinton EA. Alcohol consumption raises HDL cholesterol levels by increasing the transport rate of apolipoproteins AI and A-II. *Circulation*. 2000 Nov 7;102(19):2347-52.

10. Wakabayashi I. Cross-sectional relationship between ethanol consumption and prevalence of metabolic syndrome in Japanese men and women. *Hyogo Journal Atheroscler Thromb.* 2010.
11. Yoon YS, Oh SW, Baik HW, Park HS, Kim WY. Alcohol consumption and the metabolic syndrome in Korean adults: the 1998 Korean National Health and Nutrition Examination Survey. *The American Journal of Clinical Nutrition.* 2004.
12. Huang S, Li J, Shearer GC, Lichtenstein AH, Zheng X, Wu Y, Jin C, Wu S, Gao X. Longitudinal study of alcohol consumption and HDL concentrations: a community-based study. *The American journal of clinical nutrition.* 2017 Mar 1;105(4):905-12.
13. Vu KN, Ballantyne CM, Hoogeveen RC, Nambi V, Volcik KA, Boerwinkle E, Morrison AC. Causal role of alcohol consumption in an improved lipid profile: the Atherosclerosis Risk in Communities (ARIC) study. *PLoS One.* 2016 Feb 5;11(2):e0148765.