

To Study the Blood Pressure Response to Benidipine in Hypertension PatientsMallikarjuna Shetty*¹, Nageswar Rao Modugu², Lavanya Mandli³, Jumana Hussain⁴, Rinita Ajey⁵, Nishanth Madela⁶¹Associate Professor, Department of General medicine, M.D General medicine²Professor, Department of General medicine, M.D General medicine³Assistant Professor, Department of General medicine, M.D General medicine⁴Assistant Professor, Department of General medicine, M.D General medicine⁵Junior Resident, Department of General medicine, M.D General medicine⁶Junior Resident, Department of General medicine, M.D General medicine

The Department of General Medicine, Nizam's Institute of Medical Sciences – Hyderabad

***Corresponding author**

Mallikarjuna Shetty

Article History

Received: 10.09.2017

Accepted: 16.09.2017

Published: 30.09.2017



Abstract: Hypertension is most important cause for morbidity and mortality worldwide, most of the hypertension patients are not aware of disease and most of them are not under control. To study the clinical features, of Hypertension and blood pressure response, side effects of patients treated with Benidipine. We retrospectively collected data of all patients presenting to outpatient department over 3 months, the parameters studied were Age, symptoms, signs, investigations if any, with main focus on patients with Hypertension (Diagnosed according to JNCVII guidelines) treated with Benidipine 4mg along with blood pressure response from day1, day 8 and day 30 were noted. Total number of patients was 40 patients, with Male to Female ratio of 60%:40% and the age ranging from 34 years to 72 years with mean age of .50.72 years. The commonest symptom was headache in 55% followed by giddiness and body pains in 41.5 %, neck pain in 20 % and body pains in 17.5%, weakness in 10%. Day 1 mean Systolic Blood pressure was 162.5 mm Hg and mean Diastolic Blood pressure was 97.54 mm Hg, after treatment with Benidipine the mean Systolic Blood Pressure on day 8 was 140.6 mm Hg and mean Diastolic Blood pressure was 88.16 mm Hg, and day 30 mean systolic Blood pressure was 132.5 mm Hg and mean Diastolic Blood pressure was 86.53 mm Hg. 87.5 % patients reached target blood pressure, 12.5% did not reach the target. 2(10%) patients had pedal oedema. Benidipine has better Blood pressure lowering effects with good tolerance in hypertensive patients.

Keywords: Systolic blood pressure (SBP), Diastolic blood pressure (DBP), Hypertension, Calcium channel blocker (CCB), Benidipine.

INTRODUCTION

Hypertension is one of the main causes of mortality and morbidity globally, and major cause of cardiovascular diseases [1-3] and about 25% of rural and 33% of urban populations in India are hypertensive [4]. Among them only 10% in rural and 20% in urban have their hypertension under control [5]. Since there are many drugs used to treat hypertension, according to JNC-8 guidelines calcium channel blockers are the primary antihypertensive drugs [6].

Benidipine is a dihydropyridine-derived newer calcium channel blocker developed in Japan, with several unique mechanisms of action, that is, triple calcium channels (L, N, and T) blocking action with a membrane approach [7], and it has shown good antihypertensive effects and lesser side effects [8]. This

drug has been in India since few years and there are very few studies in Indian patients regarding efficacy and side effects which made us to collect data on Benidipine on blood pressure response and its side effects.

MATERIAL AND METHODS

Outpatient records of patients with diagnosis of Hypertension treated with Benidipine, in Nizams Institute of Medical sciences hospital which is a multispecialty, tertiary care referral hospital were collected over a period of three months.

Inclusion criteria was

- All patients diagnosed with Hypertension treated with Benidipine 4mg.

- Age above 18years. Exclusion criteria was
- Patients Age below 18 years. Information from out patient records of all patients recruited for analysis was reviewed. In the history, demographic details, symptoms with the duration, risk factors if any were noted.

Laboratory investigations

Hemogram, complete urine examination, Liver function test, renal function test, Lipid profile Chest radiograph, Ultrasound Abdomen (carried with MYLAB60 model, ESAote Company from Ahmedabad), Electrocardiography, 2D Echcardiography. Were ever done were noted.

Diagnosis

The diagnosis of Hypertension was done according to JNC VII th guidelines [9]. Hypertension was diagnosed when Systolic Blood Pressure (SBP) was >140mmHg and/or mean Diastolic Blood Pressure (DBP) > 90mmHg. Isolated systolic hypertension was defined as a Systolic Blood Pressure 140mmHg and a Diastolic Blood Pressure <90mmHg [9].

Follow up

Those patients who were on Benidipine 4mg dose, and their initial Blood pressure on day 1 was noted, then subsequently the Blood pressure was noted on day 8 and day 30 were also noted, along with any side effects.

The above data from all the patients was tabulated and analysed retrospectively with main focus on symptoms, Blood pressure response to Benidipine 4mg on day8 and day30 and any side effects were noted.

The study was retrospective audit with no patient direct identifiers; hence consent was not taken .Hospital ethics committee was informed of the study.

STATISTICAL ANALYSIS

Microsoft office 2007 was used for the statistical analysis. Descriptive statistics like mean and percentages were used to interpret the data.

RESULTS

Total 40 patients (Table-1) data was collected. with age ranging from 34 years to 72 years, with mean age being 50.72 years, Male to Female ratio of 24:16 (60%:40%). The commonest symptom being headache in 55%, next symptom was giddiness in 41.5%, neck pains in 20%, body pains in 17.5% and weakness in 10%.

Total 40 patients of Hypertension who received Benidipine 4mg (Benidin) and who came for review on day 8 and day 30 were included. The maximum Systolic blood pressure (SBP) (Table-2) noted on day1 was 194 mm of hg with mean systolic blood pressure of 162.5 mm of Hg, and maximum Diastolic blood pressure(DBP) on day 1 was 106 mm of Hg ,with mean diastolic blood pressure of 97.54 mm Hg. All the patients received Benidipine4 mg, after 8 days the mean systolic blood pressure was 140.6 mm Hg and mean diastolic blood pressure was 88.16 mm Hg. On follow up on day 30 (Figure-1) the mean systolic blood pressures was 132.5 mm Hg, and mean diastolic blood pressure was 86.53 mm Hg. That is on day 8 and day 30 Mean systolic Blood pressure came down by -21.9 mmHg and -30 mmHg respectively and mean Diastolic blood pressure came down by day 8 by -9.38 and by day 30 -11.01mm Hg from day 1 blood pressure reading. All patients reached the goal blood pressure of <140mm of Hg systolic and <90 mm Hg diastolic blood pressure, except 5 (12.5%) patients, who required Benidipine 8mg to control the blood pressure. The side effects of pedal oedema were seen in 4(10%) patients.

Table-1: shows clinical findings

	Numbers (%)
Total patients	40
Age	34 to72 years Mean age-50.72 years
Sex Male :Female	24:16 (60%:40%)
Symptoms	
Head ache	22(55%)
Giddiness	17(41.5%)
Neck pain	8(20%)
Body pains	7(17.5%)
Weakness	4 (10 %)
Side effects while on Benidipine	Pedal oedema 4 (10%)

Table-2: Showing Blood Pressure response to Benidipine 4mg


Blood pressure	Blood Pressure in mm Hg	Mean change in Blood pressure in mm Hg
Day 1		
Systolic Blood pressure	Mean-162.5	-
Diastolic Blood pressure	Mean-97.54	-
Day 8 on Benidipine 4mg		
Systolic Blood pressure	Mean-140.6	-21.9
Diastolic Blood pressure	Mean-88.16	-9.38
Day 30 on Benidipine 4 mg		
Systolic Blood pressure	Mean-132.50	-30
Diastolic Blood pressure	Mean-86.53	-11.01
Not reaching target Blood pressure by day 30	5(12.5%)	

DISCUSSION

Hypertension is one of the leading causes of coronary arterial disease and deaths from it in India [10]. In our study we had 40 patients of Hypertension who were treated with Benidipine (Benidin) 4mg. Among them 24(60%) were male and 16(40%) were females which is similarly reported by Shetty [11] *et al.* and M K Singh [12] *et al.* Of 59% to 41%, but William [13] *et al.* reported 54%to46% respectively. The age of the patients was 34 years to 72 years but M K Singh [12] *et al.* had patients age ranging from 20 years to 70 years and the mean age was 50.72 years in our study which is similarly reported by Shetty [11] *et al.* 51.72 years but William [13] *et al.* Reported mean age of 56 years in there study group.

The commonest symptom at presentation was headache in 55% followed by giddiness in 41.5%, neck pain in 20% which is similarly reported by Shetty [11] *et al.* and William [13] *et al.*

The Mean Systolic blood pressure at presentation was 162.5 mm of Hg and Mean Diastolic blood pressure was 97.54 mm of Hg which is similarly reported by Shetty [11] *et al.* but more than Gupta [14] *et al.* reported systolic blood pressure of 128.8+ - 17 mm of Hg After these patients were treated daily with Benidipine 4 mg (Benidin) the Mean Systolic Blood pressure on day 8 was 140.6 mm of Hg and Mean Diastolic pressure was 88.16 mm of Hg and on day 30 the Mean Systolic blood pressure was 132.5 mm of Hg and Mean Diastolic blood pressure

**Fig-1: Showing blood pressure response to benidipine**

Were 86.53 mm Hg, that is on day 8 and day 30 Mean Systolic blood pressure came down by -21.9 mm of Hg and -30 mm of Hg and Mean Diastolic blood pressure came down by -9.38 mm of Hg and - 11.01 mm of Hg respectively. (FIG -1).

Similarly Nakajimo [8] *et al.* reported a Mean SBP reduction from 173 mm of Hg to 135 mm of Hg and Mean DBP from 104 mm of Hg to 88 mm of Hg, but Seino [15] *et al.* reported Mean SBP decreased from 155.8 +_ 13.7 mm of Hg to 145.9 +_17 mm of Hg and Mean DBP decreased from 76.5 +_13.3mm of Hg to 71.4 +_ 13.7 mm of Hg after Benidipine 4 mg treatment

and Sasaki H *et al.* [16] showed Mean SBP decreased from 154 mm of Hg to 139 mm of Hg and Mean DBP decreased from 91 mm of Hg to 78 mm of Hg in their study.

In our study the Mean SBP reduction with Benidipine 4 mg was -30 mm of Hg and Mean DBP was -11.01 which is better than clinidipine as reported by Seino [15] *et al.* But Shetty [11] *et al.* and Bakris [17] *et al.* reported similar decrease in Mean SBP and Mean DBP with Azilasartan 40 mg. The Mean SBP and Mean DBP reduction was more with Benidipine in our study compared to candesartan reported by Rakugi [18] *et al.* of Mean SBP reduction of -17.5 mm of Hg and Mean DBP reduction of -9.8 mm of Hg. But Ihm [19] *et al.* reported similar Mean SBP and Mean DBP reduction when benidipine was compared with losartan in there study.

All the patients 87.5% reached the goal Blood Pressure of SBP of < 140 mm of Hg and DBP of < 90 mm of Hg except 12.5% patients who required Benidipine 8 mg to control Blood Pressure which is similarly reported by Nakajima [8] *et al.* and kalke s [20] *et al.* The major side effect was pedal oedema in 10 % patients.

CONCLUSION

Benidipine a new calcium channel blocker has better clinical Blood Pressure lowering effects than other calcium channel blockers, ACE inhibitors and Angiotensin receptor blockers with good tolerance. Hence it is more useful in Blood Pressure control in Hypertension patients.

Limitations

This study has very small sample size, it's a retrospective study, and does not mention about major side effects and cardiovascular mortality and about benefits over other calcium channel blockers.

ACKNOWLEDGEMENTS

To my Wife keerthi, Daugther Saanvi, Lloyed Healthcare, My patients.

REFERENCES

1. National High Blood Pressure Education Program Working Group. National High Blood Pressure Education Program Working Group Report on Hypertension in the Elderly, *Hypertension*, 1994, 23: 275–285.
2. Burt VL, Whelton P, Roccella EJ, Brown C, Cutler JA, Higgins M, Horan MJ, Labarthe D. Prevalence of hypertension in the US adult population. *Hypertension*. 1995 Mar 1;25(3):305-13.
3. O'Donnell CJ, Kannel WB. Cardiovascular risks of hypertension: Lessons from observational studies.

- Journal of Hypertension, 1998, 16 (suppl): S3–S7.7.
4. Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, Prabhakaran D. Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *Journal of hypertension*. 2014 Jun;32(6):1170.
5. Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, Prabhakaran D. Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *Journal of hypertension*. 2014 Jun;32(6):1170.
6. Takahara A. Dual L/N-Type Ca²⁺ Channel Blocker: Cilnidipine as a New Type of Antihypertensive Drug. In *Antihypertensive Drugs* 2012. InTech.
7. Yao K, Nagashima K, Miki H. Pharmacological, pharmacokinetic, and clinical properties of benidipine hydrochloride, a novel, long-acting calcium channel blocker. *Journal of pharmacological sciences*. 2006;100(4):243-61.
8. Nakajima O, Akioka H, Miyazaki M. Effect of the calcium antagonist benidipine hydrochloride on 24-h ambulatory blood pressure in patients with mild to moderate hypertension in a double-blind study against placebo. *Arzneimittelforschung*. 2000 Jul;50(07):620-5.
9. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo Jr JL, Jones DW, Materson BJ, Oparil S, Wright Jr JT, Roccella EJ. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *Jama*. 2003 May 21;289(19):2560-71.
10. Gupta R. Trends in hypertension epidemiology in India. *Journal of human hypertension*. 2004 Feb 1;18(2):73.
11. Shetty M, Nageswar RM, Lavanya M, Jummna H, Akshay P. To study the blood pressure response to azilsartan in hypertension patients. *IJCMR*, 2017;4(6):1262-1264.
12. Singh MK, Singamsetty B, Kandati J. An epidemiological study of prevalence of hypertension and its risk factors in a rural community of Nellore, Andhra Pradesh, India. *International Journal Of Community Medicine And Public Health*. 2016 Dec 22;3(12):3408-14.
13. White WB, Weber MA, Sica D, Bakris GL, Perez A, Cao C, Kupfer S. Effects of the angiotensin receptor blocker azilsartan medoxomil versus olmesartan and valsartan on ambulatory and clinic blood pressure in patients with stages 1 and 2 hypertension. *Hypertension*. 2011 Jan 1:HYPERTENSIONAHA-110.
14. Gupta R, Al-Odat NA, Gupta VP. Hypertension epidemiology in India: meta-analysis of 50 year

- prevalence rates and blood pressure trends. *J Hum Hypertens* 1996; 10: 465–472.
15. Seino H, Miyaguchi S, Yamazaki T, Ota S, Yabe R, Suzuki S. Effect of Benidipine Hydrochloride, a Long-acting T-type Calcium Channel Blocker, on Blood Pressure and Renal Function in Hypertensive Patients with Diabetes mellitus. *Arzneimittelforschung*. 2007 Aug;57(08):526-31.
 16. Sasaki H, Kanai S, Oyama T, Miyashita Y, Yamamura S, Shirai K. Effect of combination therapy of benidipine hydrochloride and candesartan cilexetil on serum lipid metabolism and blood pressure in elderly hypertensive patients with type 2 diabetes mellitus. *Journal of atherosclerosis and thrombosis*. 2006;13(3):149-57.
 17. Bakris GL, Sica D, Weber M, White WB, Roberts A, Perez A, Cao C, Kupfer S. The comparative effects of azilsartan medoxomil and olmesartan on ambulatory and clinic blood pressure. *The Journal of Clinical Hypertension*. 2011 Feb 1;13(2):81-8.
 18. Rakugi H, Enya K, Sugiura K, Ikeda Y. Comparison of the efficacy and safety of azilsartan with that of candesartan cilexetil in Japanese patients with grade I–II essential hypertension: a randomized, double-blind clinical study. *Hypertension Research*. 2012 May;35(5):552.
 19. Ihm SH, Jeon HK, Chae SC, Lim DS, Kim KS, Choi DJ, Ha JW, Kim DS, Kim KH, Cho MC, Baek SH. Benidipine has effects similar to losartan on the central blood pressure and arterial stiffness in mild to moderate essential hypertension. *Chinese medical journal*. 2012 Dec;126(11):2021-8.
 20. Kalke S, Shah BV, Nair KG, Gala D, Sood OP, Bagati A. Clinical trial of benidipine in mild to moderate hypertension. *The Journal of the Association of Physicians of India*. 1999 Feb;47(2):195-7.