

## Role of Ultrasound in Male Infertility

Dr. Vipin Raj\*

Senior Resident Department of General Surgery, SHKM GMC, NUH, Haryana, India

### Original Research Article

\*Corresponding author

Dr. Vipin Raj

#### Article History

Received: 11.04.2018

Accepted: 22.04.2018

Published: 30.04.2018

#### DOI:

10.36347/sjams.2018.v06i04.084



**Abstract:** The aims of our study is to evaluate the role of ultrasonography to identify any intra and extra testicular causes of male infertility and to correlate ultrasonographic findings with clinical and laboratory findings. This prospective study was carried out on 100 patients attending the outpatient department of Infertility clinic of Department of Surgery, detailed history and examination was done. The patients suspected to have scrotal pathology were subjected to colour Doppler sonography in the Department of Radiodiagnosis, AMU, and Aligarh. In our study clinically varicocele was the most common anomaly found followed by atrophic testes, epididymal cyst and others. On colour Doppler sonography varicocele was the most common sonographic abnormality that was present in 40 (35.08%) patients. Left sided varicocele was more common than right sided varicocele. Sonography had an overall sensitivity of 57% whereas clinical examination had an overall sensitivity of only 22%. Testicular volume and mean sperm counts/ ejaculate have shown a proportionate relationship in our study. Ultrasound is a non-invasive method, easy to perform and is cost effective. It provides important information without any post-examination consequences. Due to the high incidence of abnormalities detected by color Doppler ultrasonography we advocate that it should be used as a routine investigation in all infertile male patients.

**Keywords:** male infertility, ultrasound, varicocele.

### INTRODUCTION

It has been estimated that almost 7% of men in their reproductive age are sub-fertile or infertile due to pre-testicular, testicular or post-testicular problems Krausz [1].

Despite many technical advances that have improved diagnostic skills the etiological factors for male infertility are still obscure and around 50% of them remain undiagnosed [2]. To fill this gap in infertility clinics the use of imaging of the male genital tract has been progressively expanded. This study evaluate prospectively the role of ultrasonography in indigenous population to identify various scrotal and extra-scrotal factors with the use of scrotal ultrasound and Trans rectal ultrasound causing male infertility and there correlation with clinical and radiological findings.

### MATERIALS AND METHODS

This prospective study was carried out on 100 patients attending the outpatient department of Infertility clinic of Department of Surgery, detailed

history and examination was done. The patients suspected to have scrotal pathology were subjected to colour Doppler sonography in the Department of Radiodiagnosis, AMU, and Aligarh.

### Study subjects

Patients were selected only when two consecutive semen samples that were taken at least 5 days apart were found to be abnormal and/or if the patient had never attained fatherhood.

### Inclusion criteria

All male patients aged between 20-50 years, clinically diagnosed with infertility or sub-fertility having abnormal semen parameters as according to WHO 2010 guidelines and/or if patients had never attained fatherhood were taken for study.

**Table 1: WHO semen parameters**

Parameter	WHO 2010
Volume	1.5ml
Concentration	15 million/ml
Progressive motility	32%
Normal forms	4%

#### **Exclusion criteria**

- Patients not giving consent to be a part of the study.
- Patients having any acute scrotal pathology.
- Patients having any associated serious systemic conditions.
- Patients having incomplete medical records.
- Patients having HBsAg and HIV status positive.
- Patients who were referred to other centre after clinical, laboratory and radiological investigation.

#### **Sample size**

A total of 100 patients were taken for study that were recruited through Infertility clinic of Department of Surgery over a period of 18 months. Age of patients ranged between 20- 50 years.

#### **History and Physical Examination**

A detailed history and physical examination was done with special attention to any history suggestive of mumps, trauma to the testes, any operation and similar illness in the past. Assessment of the genitals was done with reference to examination of the penis including the location of the urethral meatus, palpation of the testes and measurement of their size, presence and consistency of the vas and epididymis, presence of varicocele, secondary sexual characteristics including body habitus, hair distribution and pattern of breast development and at last digital rectal examination.

#### **Systemic Examination**

A detailed systemic examination was done. In central nervous system assessment higher function, any neurological disorder, all cranial nerve function, sensory and motor nervous system were assessed. In cardiovascular system any abnormal heart sounds if present were documented. In respiratory system examination any illness suggesting asthma, bronchitis and any abnormalities in breath sounds were noted. In abdominal examination all hernia sites and any lump and organomegaly if present was documented

#### **Semen collection and analysis**

After 3-5 days of ejaculatory abstinence, semen samples were collected in a sterile plastic container by the process of masturbation from the subjects. Semen samples were collected in the laboratory room in a clean, dry, biologically inert container. In case of oligozoospermic or azoospermic patients, three semen samples were collected on different days with at least four days abstinence. After that gross and microscopic analysis was done and findings were noted

#### **Transrectal ultrasound**

Transrectal ultrasonography in the infertile men was done to evaluate patency of the distal ductal system (vas deferens) and internal genital organs. Sonography of all patients were done using a real time endocavitary probe (GE, 3.5 MHz). Image documentation was done with multiformat camera.

#### **Preparation and positioning and technique**

The patients were asked to take a rectal enema one night before and in the morning and instructed to take 2 to 3 glasses of water and hold urine for 2 to 4 hours before the study. The procedure was fully explained to patient in details and consent was taken.

Subjects were asked to remove their inner garments. Left lateral position was the most comfortable position for subjects in our study and this was done. Two leggings were used to cover the legs for patient's and the examiner's convenience. To make the ultrasound room more comfortable for the patient, the room was kept semi dark during assessment. The patients were explained the technique of the scanning. For transrectal ultrasound, the ultrasound transmission gel was applied to the endorectal transducer and it was covered with a sterile probe cover or a sterile condom. Scanning begins in the axial plane and the base of the prostate and seminal vesicles were visualized first. Seminal vesicles were identified bilaterally, with the ampullae of the vas on either side of the midline. The seminal vesicles are convoluted cystic structures that are darkly anechoic. Measurements were taken. Length of the seminal vesicle was taken using dotted lines along the curvature of the organ in its midline. Width was taken at the midpoint. Volume was calculated using the in-built calculator in the scanning machine. Vas deferens was visualized and the diameter of the vas deferens was measured. Next, the base of the prostate was visualized. Volume assessment of the prostate is an important and integral part of this procedure. The length and breadth of the prostate were taken on the longitudinal axis of the prostate. Thickness was taken on the transverse axis. The ellipsoid volume formula was then applied, as follows: length  $\times$  breadth  $\times$  thickness  $\times$  0.52. This was available in the in-built calculator of volume in the software on the scanning machine prostatic scanning was done in various slices.

#### **Scrotal scanning**

##### **Equipment**

A logic-500 (GE WIPRO) ultrasound machine was used. The patient was scanned using linear colour Doppler multi-frequency probe.

##### **Preparations and positioning**

For scrotal ultrasound patient is instructed to shave the parts and come for scanning. An ultrasound examination of the scrotum was done percutaneously

to investigate the testis and cord structures. The examination was done in both supine and upright position

**Technique**

The transducer (7-9 MHZ) was prepared with a sonogel (coupling gel).The transducer was held in the right hand and direct contact scan was performed with gentle rotation and angulations of the transducer. Both sagittal and transverse images were obtained. A transverse scan demonstrating both testes for comparison was obtained. Additional views were also obtained in coronal and oblique planes, with the patient upright and performing valsalva manoeuvre. Enlargement of veins was assessed. Diameter of veins was measured both in resting posture as well as in valsalva manoeuvre. Presence of varicose veins if any was noted and gradations of varicosity were done.

Colour flow Doppler examination was performed to evaluate testicular and epididymal blood

flow. The length and breadth of the testis was taken in the longitudinal axis of the testis. Thickness was taken on the transverse axis. Testicular volumes were calculated by using the empirical formula of Lambert: length  $\times$  breadth  $\times$  thickness  $\times$  0.71. This was available in the in-built calculator of volume in the software on the scanning machine. Then, the testis and para-testicular area, mediastinum testis, epididymal head, epididymal body and epididymal tail were examined sequentially. Spermatic cord and vasculature were also studied.

**STATISTICAL ANALYSIS**

The obtained data were subjected to statistical analysis using statistical software SPSS version 20.0. Data were expressed as mean  $\pm$  standard deviation and Statistical significance was analyzed by Chi-square test. Spearman's correlation test was performed to assess the linear relationship between parameters.



Scrotal probe

Endorectal probe

Fig-1: Sonography probe



**Fig-2: Sonography Apparatus**



**Fig-3: Colour Doppler sonography of scrotum in supine position**



**Fig-4: Colour Doppler Sonography of scrotum in erect position**



**Fig-5: Colour Doppler sonography of scrotum on erect position while performing valsalva manoeuvre**




**Fig-6: Transrectal sonography**

## OBSERVATIONS AND RESULTS

**Table-2: Age distribution of patients**


Serial number	Age group (in years)	Number of Patients	Percentage
1	20-24	29	29%
2	25-30	49	49%
3	31-35	10	10%
4	36-40	6	6%
5	41-45	3	3%
6	46-50	3	3%



Graph-1: Number of patients in different age groups

Table-3: Clinical findings


Serial number	Clinical findings	Number of patients (n=100)	Percentage
1	Varicocele	11	11%
2	Atrophic testis	6	6%
3	Absent testis	2	2%
4	Spermatocele/Epididymal cyst	3	3%
5	Clinically inapparent	78	78%



Graph-2: Clinical findings

Table-4: Colour Doppler sonographic findings


Serial number	Sonographic findings	Number of documentations	Percentage
1	Varicocele	40	35.08%
2	Epididymal cyst	9	7.90%
3	Atrophic testis	11	9.65%
4	Absent testis	2	1.76%
5	Testicular microlithiasis	4	3.50%
6	Bulky seminal vesicle	3	2.64%
7	Normal scan	43	37.71%
8	Prostatic cyst	2	1.76%




Graph-3: Different colour Doppler sonographic findings

Table-5: Clinical versus color Doppler findings

Serial number	Different conditions	Clinical findings	Sonographic findings
1	Varicocele	11 (11%)	40 (35.08%)
2	Epididymal cyst/Spermatocele	3 (3%)	9 (7.90%)
3	Atrophic/micro testis	6 (6%)	11 (9.65%)
4	Absent testis	2 (2%)	2 (1.76%)
5	Testicular microlithiasis	0	4 (3.50%)
6	Bulky seminal vesicle	0	3 (2.64%)
7	Prostatic cyst	0	2 (1.76%)



Graph-4: Comparison between colour Doppler and clinical examination




**Graph-5: Comparison of sensitivity of colour Doppler sonography and clinical examination**

**Table-6: Relationship between Varicocele and mean sperm count/ejaculate**

Serial number	Grades of varicocele	Number of patients	Mean sperm count/ejaculate (in millions)
1	Grade 1	13	33.15±18.7
2	Grade 2	10	34.2±28.14
3	Grade 3	19	25.37±14.9
4	Grade 4	11	15.9±8.3

**Table-7: Number of patients with different sides affected in Varicocele**


Serial number	Total number of patients	Number of patients with left sided varicocele	Number of patients with right sided varicocele
1	40	40 (40%)	21 (21%)



**Graph-6: Distribution of patients of varicocele the basis of side affection**

**Table-8: Mean sperm counts in different sonographic conditions**


Serial number	Sonographic findings	Number of documentations	Mean sperm counts/ejaculate (in millions)
1	Varicocoele	40	26.98±19.10
2	Epididymal cyst/spermatocele	9	25.11±23.87
3	Atrophic/Micro testis	11	3.45±4.845
4	Bulky seminal vesicle	3	13.0±12.12
5	Absent testis	2	4.0±5.675
6	Testicular microlithiasis	4	28.25±38.57
7	Normal scan	43	38.93±29.048



**Graph-7: Mean sperm counts among different sonographic conditions**

**Table-9: Testicular volume among infertile patients**


Serial number	Testicular volume (cc)	Number of patients (n=100)
1	0-5	6
2	6-10	6
3	11-15	26
4	16-20	28
5	21-25	20
6	26-30	8
7	31-35	1
8	36-40	5
Total		100



Graph-8: Number of patients in different testicular volume range

Table-10: Relationship between testicular volume and sperm counts


Serial number	Testicular volume (in cc)	Mean sperm counts/ejaculate (in millions)
1	0-5	1.50±3.20
2	6-10	3.83±3.746
3	11-15	17.63±12.92
4	16-20	33.43±24.951
5	21-25	42.74±25.510
6	26-30	46.63±31.55
7	31-35	88.0
8	36-40	50.4±25.245



Graph-9: Mean sperm counts in different testicular volume range

**Table-11: Relationship between Sonographic findings and testicular volume**

Serial number	Sonographic findings	Mean testicular volume (in cc)
1	Varicocele	19.20±6.58
2	Epididymal cyst/spermatocele	18.78±6.37
3	Atrophic/Micro testis	6.55±3.830
4	Bulky seminal vesicle	12±5.292
5	absent testis	4.0±2.82
6	Testicular microlithiasis	11.5±5.972
7	Normal scan	19.95±7.191
8	Prostatic cyst	10.0±7.071



**Graph-10: Mean testicular volume in different sonographic conditions**

## DISCUSSION

### Age distribution

In our study the total number of patients that were affected in the age group of 25- 30 years were 49 (49%) followed by 29 (29%) in 20-25 years, in 31-35 year there were 10 patients. Total numbers of patients in other age groups were 12 (12%) only.

In the Study conducted by Manna D, Pandit D *et al.*[3] depicted similar results in which they had showed that male infertility problems were highest in the age group of 25-34 years. In the study conducted by S. Samal, K. Dhadwe *et al.* [4] found that the maximum male infertility problems were in the age group of 31-40 years followed by other age groups. In the study conducted by Ford WC, North K *et al.* [5] concluded increased male infertility problems over the age of 40 years as compared to younger age groups, results of their study indicated that conception rate was 30 % lower in men over 40 years of age during a 12 months follow up period as compared to younger age groups. In the Study conducted by Harris I D, Randall BM *et al.* [6] stated that fertility rate in men younger than age 30 years has decreased by 15% after 1980 and also ageing of male partner has a significant adverse effect on fertility potential.

### Various sonographic conditions affecting infertile males

In our study most common clinical finding that was noted was varicocele in 11 (11%) patients followed by atrophic testes in 6 (6%) patients, epididymal cyst in 3 (3%) patients. In the study done by Sandro C Esteves, Ricardo Miyaoka *et al.*[7] revealed similar results and showed varicocele as the most common clinical anomaly in infertile patients followed by any infectious and hormonal causes In our study on colour Doppler sonography varicocele was present in 40 (35.08%) patients, epididymal cyst in 9 (7.90%) patients, atrophic testis in 11 (9.65%) patients, absent testis in 2 (1.76%) patients, testicular microlithiasis in 4 (3.50%) patients, bulky seminal vesicle in 3 (2.64%), 43 (37.71%) patients presented with a normal scan. Some of the patients in our study had two or more findings together, in those cases each finding was documented separately.

Hasan Y. Malkawi, Hussein S. Qublan *et al*<sup>8</sup> showed similar results and reported varicocele in 54% patients, epididymal cyst in 11%, testicular microlithiasis in 29% of patients, so maximum number of infertile patients in his study suffered from varicocele. Hayder M, Hayder Q. *et al.* [9] showed similar results and reported varicocele in 74%,

epididymal cyst in 8%, atrophic testes in 4%, testicular microlithiasis in 4%. This study also depicted that varicocele was the most common sonographic abnormality found in infertile male patients. Qublan HS, Al-Okoor K *et al.*[10] performed sonographic study in infertile couples and showed similar result in which he had reported varicocele in 36%, microlithiasis in 10% was the most common abnormality in infertile male patients in his study followed by others abnormalities. K.H. Tijani B.O. Oyende *et al.*[11] performed sonographic study in infertile male patients in west Africa and reported similar results and showed that varicocele was the most common abnormality in infertile couples followed by any other. P D Kantartzi, Ch D Goulis *et al.* [12]. also reported varicocele.

#### **Relationship between grades of varicocele and mean sperm counts**

In our study no definite relationship was found between grade 1 and grade 2 varicocele and mean sperm counts however, there was decrease in sperm counts in grade 3 varicocele which became more pronounced in grade 4 varicocele. In patients of grade 3 varicocele mean sperm counts were 25.37 million/ejaculate which got further deteriorated in grade 4 varicocele which was 15.9 million/ejaculate. Statistical analysis (Spearman's test) showed that negative correlation between grades 4 and 3 varicocele and mean sperm counts which was -0.81 and -0.035 respectively, which although was not statistically significant. In some patients two different grades of varicoceles were present together, in those cases each grade was documented separately.

In the study done by Cocuzza M, Athayde KS *et al*<sup>13</sup> showed similar results in which he had showed that varicocele of higher grades adversely affects semen volume and sperm counts. Belker AM[14]. also presented with similar results showed adverse effect of varicocele on overall testicular function. Pastuszak AW, Wang R[15] also concluded similar results and stated negative impact on testicular functions by varicocele, he also stated that higher grades of varicocele were associated with low sperm counts and overall poor testicular functions. Reddy SV K[16]. in his study also suggested adverse effect of varicocele on sperm counts. However study done by Hauser R *et al.*[17] stated no conclusive relationship between varicocele and sperm counts. Similarly, Jarow J[18]. in his study suggested no relationship between grades of varicocele and sperm counts.

#### **Number of patients with different sides affected in varicocele**

In our study left sided varicocele was present in 40 patients and right sided varicocele was present in 21 patients. Isolated varicocele of right side was not found in our study. Right sided varicocele was always associated with left sided varicocele.

It was due to the following reasons, the left testicular vein opens at a right angle to the left renal vein, secondly due to the fact that the loaded sigmoid colon exerts pressure on the left testicular vein and the opening of the left testicular vein is close to the opening of the adrenal veins, hence it was exposed to action of adrenergic hormones.

#### **Relationship between Testicular volume and sperm counts**

We found a positive relationship between testicular volumes and mean sperm counts. Low testicular volumes were associated with low sperm counts and as the testicular volume increases sperm count also increases.

Statistical analysis showed (Spearman's test) a highly significant positive correlation between testicular and sperm counts which was 0.617.

Similar results were showed by Arai T, Kitahara S *et al.* [19] They have also stated that testicular volume had a direct positive correlation with testicular function. Condorelli R, Calogero AE *et al*<sup>20</sup> also showed similar results and stated that low testicular volume was associated with low testicular functions. Similarly the study done by Sakamoto H[21] showed similar results and suggested that low testicular volume was associated with low testicular functions and high testicular volume was associated with higher functions. Study done by Bujan L[22] demonstrated similar results and stated that testicular volume is a key marker of testicular functions and low testicular volume was associated with low testicular functions. Kristo A[23] in his study suggested similar results and concluded a direct positive correlation between testicular volume and conventional semen parameters.

#### **Testicular volume in various sonographic conditions**

In our study testicular volume varied very abruptly in various sonographic conditions, in varicocele mean testicular volume was 19.20 cc, in epididymal cyst it was 18.78 cc, in atrophic testes it was 6.5 cc, in bulky seminal vesicle it was 12.0 cc, in absent testes it was 4.0 cc, testicular microlithiasis also presented with mean testicular volume of 11.5 cc, in prostatic cyst it was 10.0 cc, those patients that presented with normal sonographic scans had a maximum testicular volume of 19.62 cc.

These observations were possibly due to adverse effects of various conditions on testicular functions, leading to decrease in testicular volume. Although reasons for non-uniform variation of testicular volume may be due presence of two or more conditions together in some patients that produced synergistic adverse effect on testicular volume.

### Testicular volume among infertile patients

In our study maximum number of patients that attended infertility clinic had testicular volume in range of 16-20 cc, which constituted 28 patients followed by 26 patients with testicular volume in range of 11-15 cc.

In the study done by Tijani K [24] showed mean testicular volume of 15.32 cc in infertile male patients which was closely related to our study.

### Mean sperm counts in different patients

In our study there was an irregular distribution of mean sperm counts among patients with different sonographic findings. Patients of varicocele had a mean sperm counts of 26.98 million/ejaculate, in epididymal cyst mean sperm counts of 25.11 million/ejaculate, in atrophic testes mean sperm counts of 3.45 million/ejaculate. Patients that were having normal sonographic scan were having highest mean sperm counts of 35.98 million/ejaculate.

As mean sperm counts were below normal in most of these conditions. Atrophic testes affected sperm counts most adversely as already mentioned that testicular volume has a direct correlation with testicular functions. Possible causes of low sperm counts in other sonographic conditions were due to negative effects of various conditions on overall testicular functions.

Tsvetkov D [25] in his study showed similar results and stated possible negative effects of epididymal cyst on overall testicular functions leading to low sperm counts. Xu C *et al.* [26] showed similar results and stated negative effects of testicular microlithiasis on testicular functions leading to worsening of semen parameters. Contrary to our study Catanzariti F *et al.* [27] showed that testicular microlithiasis did not affect sperm parameters. In patients of absent testis Fantasia J *et al.* [28] showed similar results and stated a possible negative effect of absent testis on sperm counts

### SUMMARY

This prospective study was carried out in 100 male patients attending the outpatient department of Infertility clinic of Department of Surgery, Jawaharlal Nehru medical college and hospital over a period of 18 months.

- In our study clinically varicocele was the most common anomaly found followed by atrophic testes, epididymal cyst and others.
- On colour Doppler sonography varicocele was the most common sonographic abnormality that was present in 40 (35.08%) patients. Left sided varicocele was more common than right sided varicocele. Right sided varicocele was always associated with left sided varicocele.
- Sonography had an overall sensitivity of 57% whereas clinical examination had an overall

sensitivity of 22% only.

- There was a definite relationship between grades of varicocele and mean sperm counts. Patients that had lower grades of varicocele have a near normal sperm counts and those patients that had higher grades of varicocele were having low sperm counts.
- Testicular volume and mean sperm counts/ejaculate have shown a proportionate relationship in our study. Those patients that were having low testicular volume were having low sperm counts and as the testicular volume increases sperm counts also increases.
- We have found an uneven distribution of testicular volume among various sonographic conditions. Although testicular volume was below normal in all abnormal sonographic conditions. Patients with normal sonographic scans had largest testicular volume.
- Most sonographic abnormalities including varicocele, atrophic testes, epididymal cyst were common in 25-30 years age group. Absent testes, testicular microlithiasis and bulky seminal vesicle were not found in any specific age groups as the numbers of patients of those sonographic abnormalities were quite less.

### CONCLUSIONS

- This study emphasizes that colour Doppler ultrasonography is a very sensitive investigation in detecting various abnormalities that were often missed by repeated clinical examinations.
- Ultrasound is a non-invasive method, easy to perform and is cost effective. It provides important information without any post-examination consequences.
- Due to the high incidence of abnormalities detected by colour Doppler ultrasonography we advocate that it should be used as a routine investigation in infertile male patients.
- Given the prevalence of male infertility, the surgeon's and radiologist's familiarity with its appropriate imaging workup and recognition of the commonly involved pathologic processes is critical.
- Imaging plays a key role in the evaluation of the hypospermic or azospermic men. It can detect correctable abnormalities, which can lead to a successful conception. It can also reveal potentially life-threatening disorders in the course of an infertility evaluation.
- Ultrasonography continues to be a fast and effective modality for evaluation of the acutely painful scrotum, scrotal masses and male infertility.
- We recommend routine sonographic evaluation of all infertile male patients by USG scrotum followed by TRUS.

REFERENCES

1. Krausz C. Male infertility: Pathogenesis and clinical diagnosis. *Best Practice & Research Clinical Endocrinology & Metabolism*. 2011; 25(2):271-285
2. Tuttelmann F, Nieschlag E. Classification of Andrological disorders. In: Nieschlag E, Behre HM, Nieschlag S (eds). *Andrology. Male Reproductive Health and Dysfunction*, 3rd edn. Berlin: Springer, 2010, 87–92.
3. Manna D, Pandit D, Bhattacharya D, Biswas D. A community based study on Infertility and associated socio demographic factors in West Bengal, India. *IOSRJDMS*. 2014; 13(2):13-17.
4. Samal S, Gupta U. Epidemiological Study of Male Infertility. *Indian Medical Gazette*. 2012; Vol. CXLVI : No. 5:174-180
5. Ford WC, North K, Taylor H, Farrow A, Hull MG, Golding J. Increasing paternal age is associated with delayed conception in a large population of fertile couples: evidence for declining fecundity in older men. *Human Reproduction*. 2000 Aug 1;15(8):1703-8.
6. Harris ID, Fronczak C, Roth L, Meacham RB. Fertility and the aging male. *Reviews in urology*. 2011;13(4):e184.
7. Esteves SC, Miyaoka R, Agarwal A. An update on the clinical assessment of the infertile male. *Clinics (Sao Paulo)*. 2011;66(4):691-700
8. Malkawi HY, Qublan HS, Kakish ME, Abu-Khait SA. Frequency of scrotal abnormalities detected by ultrasound in infertile men at king Hussein medical center. *JRMS*. 2004 Jun;11(1):35-9.
9. Hayder M AL-Aridy Hayder Q. AL-Mosawi, 'The Value of Routine Scrotal Color Doppler Ultrasonography In The Infertile Male'. *QMJ* 8. (2015);12: 111-122
10. Qublan HS, Al-Okoor K, Al-Ghoweri AS, Abu-Qamar A. Sonographic spectrum of scrotal abnormalities in infertile men. *Journal of Clinical Ultrasound*. 2007 Oct 1;35(8):437-41.
11. Tijani KH, Oyende BO, Awosanya GO, Ojewola RW, Lawal AO, Yusuf AO. Scrotal abnormalities and infertility in west African men: A comparison of fertile and sub-fertile men using scrotal ultrasonography. *African Journal of Urology*. 2014 Dec 1;20(4):180-3.
12. Kantartzi PD, Goulis Ch D, Goulis GD, Papadimas I. Male infertility and varicocele: myths and reality. *Hippokratia*. 2007 Jul; 11(3):99-104
13. Cocuzza M, Athayde K, Alvarenga C, Srougi M, Hallak J. Grade 3 Varicocele in Fertile Men: A Different Entity. *The Journal of Urology*. 2012; 187(4):1363-1368
14. Belker AM. The varicocele and male infertility. *Urol Clin North Am*. 1981 Feb; 8(1):41-51.
15. Pastuszak A, Wang R. Varicocele and testicular function. *Asian Journal of Andrology*. 2015; 17(4):659.
16. Reddy SV K. Varicocele and Male Infertility: Current Issues in Management-A Review. *Medical & Surgical Urology*. 2014; 03(02).
17. Hauser R, Paz G, Botchan A, Yogev L, Yavetz H. Varicocele: effect on sperm functions. *Human Reproduction Update*, 2001, Vol.7, No.5 pp. 482-485.
18. Jarow J. Effects of varicocele on male fertility. *Human Reproduction Update*. 2001; 7(1):59-64.
19. Arai T, Kitahara S, Horiuchi S, Sumi S, Yoshida K. Relationship of testicular volume to semen profiles and serum hormone concentrations in infertile Japanese males. *Int J Fertil Womens Med*. 1998 Jan-Feb; 43(1):40-7
20. Condorelli R, Calogero A, La Vignera S. Relationship between Testicular Volume and Conventional or Nonconventional Sperm Parameters. *International Journal of Endocrinology*. 2013; 2013:1- 6.
21. Sakamoto H, Ogawa Y, Yoshida H. Relationship between testicular volume and testicular function: comparison of the Prader orchidometric and ultrasonographic measurements in patients with infertility. *Asian J Androl*. 2008 Mar;10(2):319-24
22. Bujan L, Mieusset R, Mansat A, Moatti JP, Mondinat C, Pontonnier F. Testicular size in infertile men: relationship to semen characteristics and hormonal blood levels. *Br J Urol*. 1989 Dec;64(6):632-7.
23. Kristo A, Dani E. The Correlation between Ultrasound Testicular Volume and Conventional Semen Parameters in Albanian Subfertile Males. *Maced J Med Sci*. 2014
24. Tijani K, Oyende B, Awosanya G, Ojewola R, Yusuf A. Assessment of testicular volume: A comparison of fertile and sub- fertile West African men. *African Journal of Urology*. 2014; 20(3):136-140
25. Tsvetkov D. [Epididymal cysts, an etiological factor for male infertility]. *Akush Ginekol (Sofia)*. 1987;26(2):53-7.
26. Xu C, Liu M, Zhang FF, Liu JL, Jiang XZ, Teng JB, Xuan XJ, Ma JL. The association between testicular microlithiasis and semen parameters in Chinese adult men with fertility intention: experience of 226 cases. *Urology*. 2014 Oct;84(4):815-20
27. Catanzariti F, Cantoro U, Lacetera V, Muzzonigro G, Polito M. Testicular microlithiasis and dyspermia: Is there any correlation?. *Archivio Italiano di Urologia e Andrologia*. 2014 Mar 28;86(1):20-2.
28. Fantasia J, Aidlen J, Lathrop W, Ellsworth P. Undescended Testes: A Clinical and Surgical Review. *Urol Nurs*. 2015 May-Jun;35(3):117-26.