

## Research Article

# Incidence of Recurrent Laryngeal Nerves Injury during Thyroid Surgery

Saadeldin A. Idris<sup>1\*</sup>, Qurashi M. Ali<sup>2</sup>, Aamir A. Hamza<sup>3</sup>

<sup>1</sup>Associate professor, Alzaeim Alazhari University, Faculty of Medicine, Khartoum, Sudan.

<sup>2</sup>Professor of anatomy, National College for Medical and Technical Studies, Khartoum, Sudan.

<sup>3</sup>Associate professor, Bahri University, Faculty of Medicine, Khartoum, Sudan.

### \*Corresponding author

Saadeldin A. Idris

Email: [saadeldinahmed@hotmail.com](mailto:saadeldinahmed@hotmail.com)

**Abstract:** The technique of thyroidectomy has been in evolution for many years. It is a fundamental principle of surgery that a structure must be identified clearly during the procedure in order to prevent its damage. The objective of the study was to evaluate our routine identification of the recurrent laryngeal nerve (RLN) during thyroidectomy aiming to lessen the inadvertent injury of the recurrent laryngeal nerves during surgery. The study included a convenience sample, non probability total coverage multicentre hospital based descriptive prospective study (December 2009 to May 2012) including all adult patients managed by subtotal thyroidectomy after acceptance of the pre given informed consent. All operations were undertaken by the same surgical team. Data collected using a predesigned questionnaire. Data analysis was performed using SPSS version 15.0 for Windows. The study included 82 patients (69 (84.15%) females and 13(15.85%) males), with male to female ratio of 1:5.3. Their mean age was 42.8 years (SD ± 8.4 years). All were treated by subtotal thyroidectomy. The rate of identification of the RLN was 100% for all patients. The overall frequency of RLN injury was 1.2% seen in one patient. It was temporary injury. In conclusion, study proved that surgical exposure of the RLNs avoids the incidence of its damage.

**Keywords:** Thyroidectomy, Recurrent laryngeal nerve, identification, injury.

## INTRODUCTION

Thyroidectomy is globally practiced to treat a wide range of thyroid swellings and is considered as a safe procedure in well equipped settings with suitable experience to anticipate and avoid the occurrence of possible surgical complications [1].

Complications of any surgical procedure are a sensitive measure of the quality. Reported complications following thyroid surgery are rare but their consequences can often be life -threatening as compared to the some other surgeries being performed routinely [1, 2].

One of the most feared complications of thyroid surgery is the Recurrent Laryngeal Nerve (RLN) injury [3, 4], also the most common reason of litigation related to the complications of thyroid surgery [5]. RLN injury is an annoying but avoidable complication which results from severing, clamping or stretching of the nerve during surgery and may result in severe untoward sequelae for the patient [6].

Review of literature revealed that the prevalence of RLN palsy varies from centre to centre depending upon the level of experience in thyroid surgery and the nature of surgery [1, 2, 6, 7]. The exact incidence of recurrent laryngeal nerve injury is unknown and varies widely [6, 8]. There is controversy in whether the identification of the recurrent laryngeal nerve during the procedure will affect the incidence of its damage or not [9].

In an attempt to improve our results of thyroidectomy we have undertaken a careful study using RLN visualization technique to determine the incidence of RLN injury in our practice.

## PATIENTS AND METHODS

A convenience sample, non probability total coverage multicentre Hospital based descriptive prospective study was carried out in Charity teaching hospital and Omdurman teaching hospital between December 2009 and May 2012. All adult patients from both genders with thyroid disorder that were managed surgically by subtotal thyroidectomy were enrolled into study after being accepted the pre given informed consent.

Patients with previous thyroid surgery or with malignant thyroid disease were excluded from the study. Preoperative diagnosis and status of thyroid disease was done clinically, thyroid function tests and histopathologically by fine needle aspiration or Tru cut needle biopsy when needed.

Our routine operative strategy to thyroid glands and the surgical technique that we have adopted has been described and standardized earlier [10].

Subtotal thyroidectomy is carried out through a transverse 'collar' incision, two fingers' breadth above the suprasternal notch. This lies in the line of the natural

skin folds of the neck. Skin flaps are reflected, together with platysma, and the investing fascia opened longitudinally between the strap muscles and between the anterior jugular veins. If more room is required in the case of a large goitre, the strap muscles are divided; this is carried out at their upper extremity because their nerve supply (the ansa hypoglossi) enters the lower part of the muscles and is hence preserved.

The pretracheal fascia is then divided, exposing the thyroid gland. The thyroid is then mobilized and its vessels ligated after identification of the RLN during the ligation of the inferior thyroid artery (ITA) and superior laryngeal nerve during the ligation of the superior thyroid artery in each side separately. All operations were undertaken in both hospitals by the same surgical team.

Data collected using a predesigned questionnaire. All thyroidectomies performed during the period of study were evaluated, and the various types of recurrent laryngeal nerves were recorded. Data analysis was performed using SPSS version 15.0 for Windows.

**RESULTS**

The study included 82 patients (69 (84.15%) females and 13(15.85%) males), with male to female ratio of 1:5.3. Their mean age was 42.8 years (SD±8.4 years) (Table 1). The indication of thyroidectomy was variable as indicated in table 2.

**Table 1: Patients’ characteristics (n=82)**


Patients’ characteristics	n = 82
Age (mean ± SD)	42.8 ± 8.4
Gender (♂/♀)	13/69
Type of surgery	
Subtotal thyroidectomy	82
Permanent recurrent laryngeal nerve palsy	0
Transient recurrent laryngeal nerve palsy	1

**Table 2: Classification of thyroid disorders (n=82)**

Indication for surgery	Frequency	Percent
Diffuse goitre with obstructive symptoms	40	48.8
Multinodular goitre	31	37.8
Dominant Single large nodular goitre	11	13.4
Total	82	100.0

All were treated by subtotal thyroidectomy (bilateral resection with leaving a slice of thyroid tissue on each side). The rate of identification of the RLN was 100% for all patients (Fig. 1).

Variations such as a non-recurrent laryngeal nerve or the absence of the ITA were not found in this study.



**Fig.1: Surgical exposure of the recurrent nerve (yellow arrow the recurrent laryngeal nerve, red arrow inferior thyroid artery (ITA)) (n=82)**

The overall frequency of RLN injury was unilateral right sided and found to be 1.2% seen in one patient. It was confirmed by direct laryngoscope. It was temporary injury and recovered full function within 4 weeks postoperatively.

**DISCUSSION**

Thyroidectomy is the surgery in which RLN injury occurs most frequently. Its injury is an unwelcome and not infrequent complication during neck dissection [7, 11, 12]. Surgically induced recurrent laryngeal nerve paralyses are frequently not recognized at the time of thyroid surgery [7].

According to our observation in the current study, the overall frequency of RLN injury was found to be 1.2% seen in one patient. It was temporary injury. The injury was noticed immediately after surgery, when on first post operative day patient had hoarseness of voice. This patient had undergone subtotal thyroidectomy for multinodular goitre. Mishra et al, Thomusch et al, Steurer et al, Wagner et al, and Deus et al at their studies showed that the incidence of recurrent laryngeal nerve injuries ranging between 0.0%-13% [3, 13-16]. Others reported incidence of permanent injury to RLN ranges from 0% to 5.2% and for temporary injury from 0.4% to 7.2% [3, 4, 8].

According to Titche [17], thyroidectomy accounts for 35.71% of surgical causes of injury to the RLN, and accounts for 3.73% of all causes.

Saadeldin A. Idris et al. [18] in the study of outcomes and complication of thyroid surgery among the Sudanese patients, during 1351 thyroidectomies observed incidence of recurrent laryngeal nerve injury was 1.9% (26 cases). Of which, the incidence of transient unilateral RLN paralysis was 1.2% (16 cases), permanent unilateral RLN palsy was 0.5% (7 cases),

transient bilateral RLN palsy was 0.2% (3 cases), and no permanent bilateral RLN paralysis detected.

Several studies revealed that depending upon the skill of an individual surgeon principal identification of the nerve reduces the risk of permanent laryngeal nerve injuries from over 5% to less than 1% [16, 19-23]. Wagner and Seiler [16] in their study of recurrent laryngeal nerve palsy after thyroid gland surgery in 1026 patients, the incidence of transient and permanent RLN was 5.9% and 2.4% respectively. Jatzko *et al.* [19] in a series of 803 patients the incidence of transient RLN palsy was 3.6% and for permanent RLN palsy it was 0.5%. Sosa *et al.* [20] in their large study of 5860 patients, they found that the incidence of permanent RLN palsy 0.8 % and they did not recorded any transient RLN palsy. Hermann *et al.* [21] in their study of laryngeal recurrent nerve injury in surgery for benign thyroid diseases: effect of nerve dissection and impact of individual surgeon in nerves at risk, study analyzed 9,385 patients, the incidence of permanent RLN palsy was 3.0%. Rosato *et al.* [22] in their longitudinal analysis of a multicentric study of 14,934 patients, the incidence of transient and permanent RLN palsy was 2.0% and 1.0% respectively. Goncalves and Kowalski [23] in study of surgical complications after thyroid surgery in 1020 patients, the incidence nerve palsy was 1.4% for transient RLN palsy and 0.4% for permanent RLN palsy.

Several factors influence the likelihood of injury to the nerve, including the underlying disease, the extent of resection, and the experience of the surgeon. Bleeding should and can be kept to a minimum and the use of diathermy should be avoided in the vicinity of the laryngeal nerves [24].

Rosato and other authors have reported on various factors that could lead to voice changes post-thyroidectomy besides iatrogenic injury [24-26]. This underlines the importance of both pre and post operative voice assessment, although Yeung has found indirect laryngoscopy of limited value in the preoperative assessment of symptomatic patients [27].

## CONCLUSION

The study proved earlier reports that surgical exposure of the RLNs avoids the incidence of its damage. We advocate the routine identification and dissection of RLN to reduce its injury risk to minimum.

## REFERENCES

1. Doumi EA, Mohamed IM, Abakar AM, Bakhiet MY; Thyroidectomy at El Obeid Hospital, Western Sudan. *Khartoum Medical Journal*, 2009; 2(1):158-161.
2. Prim MP, De Diego JI, Hardisson D, Madero R, Gavilan J; Factors related to nerve injury and hypocalcemia in thyroid gland surgery.

- Otolaryngol Head Neck Surg., 2001; 124(1): 111-114.
3. Steurer M, Passler C, Denk DM, Schneider B, Niederle, Bigenzahn W; Advantages of recurrent laryngeal nerve identification in thyroidectomy and parathyroidectomy and the importance of preoperative and postoperative laryngologic examination in more than 1000 nerves at risk. *Laryngoscope*, 2002; 112(1):124-133.
4. Chiang FY, Wang LF, Huang YF, Lee KW, Kuo WR; Recurrent laryngeal nerve palsy after thyroidectomy with routine identification of recurrent laryngeal nerve. *Surgery*, 2005; 137(3): 342-347.
5. Ready AR, Barnes AD; Complications of thyroidectomy. *Br J Surg.*, 1994; 81:1555-1556.
6. Souza LS, Crespo AN, Alves de Medeiros JL; Laryngeal vocal and endoscopic alterations after thyroidectomy under local anesthesia and hypnosis. *Braz. J. Otorhinolaryngol. (Impr.)*, 2009; 75(4): 213-220.
7. Myssiorek D; Recurrent Laryngeal nerve paralysis: anatomy and etiology. *Otolaryngol Clin N Am.*, 2004; 37(1): 25-44.
8. Ardito G, Revelli R, d'Alatri L, Lerro V, Guidi ML, Ardito F; Revisited anatomy of the recurrent laryngeal nerves. *Am J Surg.*, 2004; 187(2): 249-253.
9. Hisham AN, Lukman MR; Recurrent laryngeal nerve in thyroid surgery: A critical appraisal. *ANZ J Surg.*, 2002; 72(12): 887-889.
10. Riddell VH; Injury to the recurrent laryngeal nerves during thyroidectomy ; a comparison between the results of identification and non-identification in 1022 nerves exposed to risk.. *Lancet*, 1956; 271 (6944): 638-641.
11. Liebermann-Meffert DM, Walbrun B, Hiebert CA, Siewert JR; Recurrent and superior laryngeal nerves: a new look with implications for the esophageal surgeon. *Ann Thorac Surg.*, 1999; 67(1): 217-223.
12. Mc Henry CR; Patient volumes and complications in thyroid surgery. *Br J Surg.*, 2002; 89(7): 821-823.
13. Mishra A, Agarwal A, Agarwal G, Mishra SK; Total thyroidectomy for benign thyroid disorders in an endemic region. *World J Surg.*, 2001; 25(3): 307-310.
14. Thomusch O, Machens A, Sekulla C, Ukkat J, Lippert H, Gastinger I *et al.*; Multivariate analysis of risk factors for postoperative complications in benign goiter surgery: prospective multicenter study in Germany. *World J Surg.* 2000; 24(11):1335-1341.
15. Deus Fombellida J, Gil Romea I, García Algara C, Sancho MA, Alonso Gotor MA, Morino Mirallas MJ *et al.*; Aspectos quirúrgicos de los bocios multinodulares. *A*

- propósito de una serie de 680 casos. *Cir Esp.* 2001; 69(1): 25–29.
16. Wagner HE, Seiler C; Recurrent laryngeal nerve palsy after thyroid gland surgery. *Br J Surg.*, 1994; 81(2): 226–228.
  17. Titcher LL; Causes of recurrent laryngeal nerve paralysis. *Arch Otolaryngol.*, 1976; 102(5): 259-261.
  18. Idris SA, Shalayel MH, Idris A, Ali AQ; Outcomes and complications of thyroid surgery among the Sudanese patients. *Sudan Medical Monitor*, 2008; 3(4):143-148.
  19. Jatzko GR, Lisborg PH, Müller MG, Wette VM; Recurrent nerve palsy after thyroid operations: principal nerve identification and a literature review. *Surgery*, 1994; 115(2):139–144.
  20. Sosa JA, Bowman HM, Tielsch JM, Powe NR, Gordon TA, Udelsman R; The importance of surgeon experience for clinical and economic outcomes from thyroidectomy. *Ann Surg.*, 1998; 228(3): 320–330.
  21. Hermann M, Alk G, Roka R, Glaser K, Freissmuth M; Laryngeal recurrent nerve injury in surgery for benign thyroid diseases: effect of nerve dissection and impact of individual surgeon in more than 27,000 nerves at risk. *Ann Surg.*, 2002; 235(2): 261–268.
  22. Rosato L, Avenia N, Bernante P, De Palma M, Gulino G, Nasi PG *et al.*; Complications of thyroid surgery: analysis of a multicentric study on 14,934 patients operated on in Italy over 5 years. *World J Surg.*, 2004; 28(3): 271–276.
  23. Goncalves Filho J, Kowalski LP; Surgical complications after thyroid surgery performed in a cancer hospital. *Otolaryngol Head Neck Surg.*, 2005; 132(3): 490–494.
  24. Rosato L, Carlevato MT, De TG, Avenia N, Rosato L, Carlevato MT *et al.*; Recurrent laryngeal nerve damage and phonetic modifications after total thyroidectomy: surgical malpractice only or predictable sequence? *World Journal of Surgery*, 2005; 29(6): 780-784.
  25. Crookes PF, Recabaren JA, Crookes PF, Recabaren JA; Injury to the superior laryngeal branch of the vagus during thyroidectomy: lesson or myth? *Annals of Surgery*, 2001; 233(4): 588-593.
  26. Sinagra DL, Montesinos MR, Tacchi VA, Moreno JC, Falco JE, Mezzadri NA *et al.*; Voice changes after thyroidectomy without recurrent laryngeal nerve injury. *Journal of the American College of Surgeons*, 2004; 199(4): 556-560.
  27. Yeung P, Erskine C, Mathews P, Crowe PJ, Yeung P, Erskine C *et al.*; Voice changes and thyroid surgery: is pre-operative indirect laryngoscopy necessary? *Australian & New Zealand Journal of Surgery*, 1999; 69(9): 632-634.