

Research Article**Blood Group Distribution in a Study Population and their Associated Rhesus Factor (Uganda)****Frank Kamya Mukinda^{1*}, Herman Lule¹, Muhamudu Kalange², Isaac Echoru³, Herbert Izo Ninsiima², Keneth Iceland Kasozi²**¹Department of Surgery, School of Clinical Medicine and Dentistry, Kampala International University Western Campus Teaching Hospital, Bushenyi, Uganda²Department of Physiology, Faculty of Biomedical Sciences, Kampala International University Western Campus, Bushenyi, Uganda³Department of Anatomy, Faculty of Biomedical Sciences, Kampala International University Western Campus, Bushenyi, Uganda***Corresponding author**

Frank Kamya Mukinda & Keneth Iceland Kasozi

Email: rockfrank17@yahoo.com and kicelandy@gmail.com

Abstract: Whole blood transfusions have continuously been done in medical hospitals as the major line of emergency recovery on the African continent as a result of limited infrastructure despite of the fact that the general blood group status is poorly understood. The aim of the study was to determine the prevalent blood groups and their associated rhesus factor. This was a cross sectional study conducted for a period of two months at Kampala International University Teaching Hospital involving a total of 47 undergraduate students, blood was picked by pricking the index finger and placed on a white tile from which the antigens were added to it. The study revealed that majority (53.2%) of the population had blood group O with the greatest prevalence being in the male population and blood group B was the second prevalent blood group in this study. This would probably be due to the establishment of endemic stability with malaria infections in the community. This would be due to genetic selection thus favoring the proliferation of the B and O blood groups as they appear to offer a mutuality protective advantage against plasmodium parasites. The major associated Rhesus factor were of O⁺ (53.2%) and B⁺ (21.3%) which showed that there was a significant downward shift in the levels of the Rhesus antigen. Blood group O was the most prevalent probably due to its evolutionary advantage in the region. Phenotypic expression need to be investigated further for the development of a strong hematological team in various rural communities.**Keywords:** Blood groups, Uganda, O prevalence

INTRODUCTION

In sub Saharan countries like Uganda, majority of blood transfusions are carried out using the routine blood typing protocols [1]. Blood transfusions have been shown to be associated with enormous health risks and these have been shown to either be immediate or delayed thus making clinical diagnosis and reporting of cases different to monitor [2]. Epidemiological information on the hematological status of sub Saharan patients is important for a full scale management of hematological related cases in the communities. This requires the establishment of reliable hematological reports that offer fundamental knowledge to policy makers in the region in preparation for future projects. The major internationally recognized blood group systems in operation are the ABO, Rhesus (Rhesus) systems [3, 4]. Recent funding in various regional hospitals through the World Health Organization (WHO) has led to an improvement of blood handling

leading to a reduction in unnecessary transfusion accidents [5]. The increasing prevalence of infectious and non-infectious diseases ranging from HIV/AIDS, hepatitis, malaria to the hemolytic disease of the new born, as well as chronic cardio-pulmonary challenges has led to increased research in the field of molecular hematology [3, 6-10]. Studies have shown the distribution of the ABO phenotypes to vary among different populations [3, 4]. The A phenotype is the most prevalent in North Central Europe while the B phenotype is most frequent in Central Asia [3, 11]. Blood group O is the most prevalent phenotype globally, with majority of Africa and Australia showing highest frequencies [12]. It's thought to be due to evolutionary selection based on pathogen-driven blood group antigen changes. The current status of the prevalent blood groups in sub Saharan Uganda is poorly known, thus the study was conducted to determine the recent blood frequencies in Western Uganda.

MATERIALS AND METHODS

This was a cross sectional study conducted for a period of two months (October – November 2014) at Kampala International University Teaching Hospital involving a total of 47 undergraduate students. These were randomly chosen, grouped into two groups basing on sex and their blood groups were determined using commercial anti-sera. Briefly, 2 ml of venous blood were collected into a sterile vacutainer containing EDTA. 2 drops of the blood were applied on a clean white tile and an equal value of anti-serum added to determine the blood group and Rhesus factors.

Statistical analysis

Data was recorded as frequencies and expressed as percentages. Group comparisons on gender were carried out using SPSS Version 21 and $p < 0.05$ was considered statistically significant.

Ethical Considerations

All authors hereby declare that all experiments have been examined and approved by then appropriate ethics committee and have therefore been performed

in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

Consent

Informed consent was obtained from participants before they were enrolled for this study.

RESULTS

The study revealed that majority (53.2%) of the population had blood group O with the greatest prevalence being in the male population. Blood group B was also found to be highly prevalent (23.4%) and second to blood group O in which the male population were also most prevalent (14.9%). The positive Rhesus factor was shown to be the most prevalent (89.4%) antigen in the population and it was most prevalent in the male population (61.7%) as shown in Table 1. No statistical significance ($p = 0.486$) was shown to exist between gender and the different blood groups. The Rhesus factor prevalence in the participants was also found to be independent ($p = 0.613$) of sex.

Table 1: Showing frequencies of blood groups among participants

Parameter	Variable	Frequency of gender of participants (%)		
		Female	Male	Total
Blood group	A	0 (0)	5 (10.6)	5 (10.6)
	B	4 (8.5)	7 (14.9)	11 (23.4)
	AB	2 (4.3)	4 (8.5)	6 (12.8)
	O	8 (17)	17 (36.2)	25 (53.2)
Rhesus factor	Positive	13 (27.7)	29 (61.7)	42 (89.4)
	Negative	1 (2.1)	4 (8.5)	5 (10.6)

The Rhesus factor was found to be most prevalent (53.2%) in blood group O with the positive (O^+) Rhesus factor being most prevalent (48.9%). This was followed by blood group B (23.4%) and the most

prevalent genotype was of the positive factor (B^+) at 21.3% as shown in Table 2. No statistical significance ($p = 0.825$) was shown between Rhesus factor and blood group in the population.

Table 2: Showing distribution of Rhesus factor amongst blood groups

Blood group	Rhesus factor frequency (%)		
	Positive	Negative	Total
A	4 (8.5)	1 (2.1)	5 (10.6)
B	10 (21.3)	1 (2.1)	11 (23.4)
AB	5 (10.6)	1 (2.1)	6 (12.8)
O	23 (48.9)	2 (4.3)	25 (53.2)

DISCUSSION

The study revealed that majority (53.2%) of the population had blood group O with the greatest prevalence being in the male population as shown in Table 1. This would probably be due to the small sample size which had a male to female ratio of 3:1. Blood group O has been shown to be the most prevalent in the black population in previous study [13, 14] and blacks of American decent have been shown to be associated with a sickle cell trait (SCT) thus showing that genotypic expression are major factors in the determination of pathological outcomes of disease [15]. Since the Rhesus factor is located on the surface of the

RBCs, presence or absence of the factor depends on the genetic distribution of the genes during meiosis and since the entire process is random, one's sex doesn't necessary have any association ($p = 0.613$) with the type of Rhesus factor which will be expressed. Blood group B was the second prevalent blood group in this study which would probably due to the establishment of endemic stability with malaria infections in the community. This would be due to genetic selection thus favoring the proliferation of the B and O blood groups as they appear to offer a protective advantage against plasmodium parasites and this has been shown to have

serious implications for blood donation campaigns [4, 16, 17].

The study further revealed that the Rhesus factor was most prevalent (53.2%) in blood group O i.e. O⁺ followed by B⁺ (21.3%) as shown in Table 2. These findings show a significant downward shift in the levels of the Rhesus antigen compared to a recent study in West Africa [1]. This would probably be due to differences in study area, populations size, genotypic variation, and environmental evolutionary changes [1, 4, 11]. No statistical significance ($p = 0.825$) was shown between Rhesus factor and blood group in the population. This is due to the fact that Rhesus factor is determined by genetic factors which are in built within the DNA of the RBCs [4].

CONCLUSION AND RECOMMENDATIONS

In conclusion, the most prevalent blood group types were O⁺ followed by B⁺ which is thought to be due to the evolutionary advantage the groups have against the endemic plasmodium parasites in the region. Further studies would be carried out to determine the phenotypic expression and genetic predisposition to plasmodium parasites in the study community for the development of a strong hematological team in various rural communities. This would help to provide precise and accurate knowledge on the management of patients of East sub Sahara African origin, especially in the rural communities which struggle with minimal infrastructure.

ACKNOWLEDGEMENT

The authors would like to thank all participants that participated and helped to make this work a success. All persons in the teaching hospital of Kampala International University Western Campus who participated in the study especially during the data collection process are henceforth acknowledged.

REFERENCES

1. Bogui LS, Dembele B, Sekongo Y, Abisse S, Konaté S, Sombo M; Phenotypic Profile of Rh and Kell Blood Group Systems among Blood Donors in Cote d ' Ivoire , West Africa. *Journal of Blood Transfusion*, 2014; 2014, Article ID 309817: 4 pages. Available from <http://www.hindawi.com/journals/jbt/2014/309817/>
2. Fleming AF; HIV and blood transfusion in sub-Saharan Africa. *Transfusion science*, 1997; 18(2): 167-179.
3. Arifuzzaman M, Ahmed T, Rahman MA, Chowdhury F, Rashu R, Khan AI *et al.*; Individuals with Le (a + b 2) Blood Group Have Increased Susceptibility to Symptomatic *Vibrio cholerae* O1 Infection, *PLoS Negl Trop Dis.*, 2011; 5(12): e1413.
4. Hansson HH, Kurtzhals JA, Goka BQ, Rodrigues OP, Nkrumah FN, Theander TG *et al.*; Human genetic polymorphisms in the Knops blood group are not associated with a protective advantage against *Plasmodium falciparum* malaria in Southern Ghana. *Malar J.*, 2013; 12: 400.
5. Osaro E, Charles AT; The challenges of meeting the blood transfusion requirements in Sub-Saharan Africa: the need for the development of alternatives to allogenic blood. *J Blood Med.*, 2011; 2: 7–21.
6. Fleming AF; AIDS in Africa--an update. *AIDS Forsch.*, 1988; 3(3): 116–138.
7. Mensah GA; Epidemiology of stroke and high blood pressure in Africa. *Heart (British Cardiac Society)*, 2008; 94(6): 697–705.
8. Olsen MA, Butler AM, Willers DM, Gross GA, Devkota P, Fraser VJ; Risk factors for endometritis after low transverse cesarean delivery. *Infection Control and Hospital Epidemiology: The Official Journal of the Society of Hospital Epidemiologists of America*, 2010; 31(1): 69–77.
9. Jones CE, Hesseling AC, Tena-Coki NG, Scriba TJ, Chegou NN, Kidd M *et al.*; The impact of HIV exposure and maternal Mycobacterium tuberculosis infection on infant immune responses to bacille Calmette-Guérin vaccination. *AIDS (London, England)*, 2015; 29(2): 155–165.
10. Cserti-Gazdewich CM, Dhabangi A, Musoke C, Ssewanyana I, Ddungu H, Nakiboneka-Ssenabulya D *et al.*; Cytoadherence in paediatric malaria: ABO blood group, CD36, and ICAM1 expression and severe *Plasmodium falciparum* infection. *Br J Haematol.*, 2012; 159(2): 223–236.
11. Barragan A, Kremsner PG, Wahlgren M, Carlson J; Blood Group A antigen is a coreceptor in *Plasmodium falciparum* rosetting. *Infection and Immunity*, 2000; 68(5): 2971–2975.
12. Zhang H, Mooney J, Reilly MP; ABO blood groups and cardiovascular diseases. *International Journal of Vascular Medicine*, 2012; 2012, Article ID 641917: 11 pages. Available from <http://www.hindawi.com/journals/ijvm/2012/641917/>
13. Peeters M, Gueye A, Mboup S, Bibollet-Ruche F, Ekaza E, Mulanga C *et al.*; Geographical distribution of HIV-1 group O viruses in Africa. *AIDS (London, England)*, 1997; 11(4): 493–498.
14. Fang C, Cohen HW, Billett HH; Race, ABO blood type and VTE Risk: Not black and white. *Transfusion*, 2013; 53(1): 187–192.
15. Yazdanbakhsh K, Ware RE, Noizat-Pirenne F; Red blood cell alloimmunization in sickle cell disease: pathophysiology, risk factors, and transfusion management. *Blood*, 2012; 120(3): 528–537.
16. Panda AK, Panda SK, Sahu AN, Tripathy R, Ravindran B, Das BK; Association of ABO blood group with severe falciparum malaria in adults : case control study and meta-analysis. *Malaria Journal*, 2011; 10(1): 309.
17. Tagny CT, Owusu-Ofori S, Mbanya D, Deneys V; The blood donor in sub-Saharan Africa: a review. *Transfus Med.*, 2010; 20(1):1-10.