

**Research Article****Cytomorphological Patterns of Tuberculous Lymphadenitis: Experience from a Tertiary Centre in Rural Haryana**Shweta Rana<sup>1</sup>, Puja Sharma<sup>2</sup>, Shivani Kalhan<sup>3</sup>, Pawan Singh<sup>4</sup>, Manmeet Kaur Gill<sup>5</sup>, Arun Kumar<sup>6</sup><sup>1</sup>Demonstrator, Dept. of Pathology, SHKM, GMC, Nalhar, Mewat, Haryana.<sup>2</sup>Associate Professor, Dept. of Pathology, SHKM, GMC, Nalhar, Mewat, Haryana.<sup>3</sup>Professor and Head, Dept. of Pathology, SHKM, GMC, Nalhar, Mewat, Haryana.<sup>4</sup>Assistant Professor, Dept. of Pathology, SHKM, GMC, Nalhar, Mewat, Haryana.<sup>5</sup>Demonstrator, Dept. of Pathology, SHKM, GMC, Nalhar, Mewat, Haryana.<sup>6</sup>Assistant Professor, Dept. of Community Medicine, SHKM, GMC, Nalhar, Mewat, Haryana.**\*Corresponding author**

Dr. Shweta Rana

Email: [drshwetarana15@gmail.com](mailto:drshwetarana15@gmail.com)

---

**Abstract:** Tuberculous lymphadenitis is the commonest cause of lymphadenopathy in developing countries. Cytomorphology along with acid fast staining plays an important role in diagnosing these cases, thus avoiding unnecessary surgical intervention. The aim is to study the epidemiological patterns and cytomorphological presentations of tuberculous lymphadenitis in reference to Ziehl- Neelsen staining. This was a retrospective study and a total of 891 patients including all age groups and both sexes presenting with palpable or deep lymph nodes in fine needle aspiration cytology (FNAC) clinic of our institute over a period of 2 years and four months were included in the study. Smears were fixed in 95% ethyl alcohol and stained with Papanicolaou stain. Leishman stain was done on air dried smears. Ziehl-Neelsen(ZN) staining was done wherever required. During this period, 500 cases of Tuberculous lymphadenitis were diagnosed on cytology. These were further categorised into three patterns based on cytomorphological features and the percentage of acid fast bacilli (AFB) positive cases in each category were noted. Data was entered in excel spread sheet and statistical analysis was done. P value <0.05 was taken to represent significant difference. Out of 891 aspirations from lymph nodes, 500 cases were diagnosed as Tuberculous lymphadenitis. Three cytomorphological patterns were observed. These were- Group 1: Granulomas without necrosis (26.6%), Group 2: Caseating epithelioid Granulomas (66%), Group 3: Necrotizing lymphadenitis (7.4%). Overall, AFB positivity was seen in 21.6% cases. The incidence of Tuberculous lymphadenitis in our study population is significantly high. FNAC is useful in diagnosing Tuberculous lymphadenitis and its sensitivity can further be increased by complementing cytomorphology with acid fast staining.**Keywords:** FNAC, Tuberculous lymphadenitis, Epithelioid cell granuloma, Cytomorphological patterns, Necrotizing lymphadenitis, Extrapulmonary tuberculosis.

---

**INTRODUCTION**


Tuberculous lymphadenitis is the most common form of extrapulmonary tuberculosis [1]. In developing countries, almost two third of the cases of lymphadenopathy are due to tuberculosis [2]. FNAC as first line of investigation has assumed importance in diagnosing a variety of disease processes as it is rapid, simple, reliable, minimally invasive and cost effective procedure which can be used in outpatient setting [3]. FNAC is economical and rapid as compared to culture studies (considered as gold standard but time consuming) and Polymerase chain reaction; which is expensive [4]. Tuberculosis is very common in our country and tuberculous lymphadenitis is the most common cause of chronic lymphnode enlargement [5]. Though the most reliable criteria for diagnosing tuberculous lymphadenitis is demonstration of acid fast

bacilli (AFB) by Ziehl- Neelsen (ZN) stain, auramine rhodamine stain or by polymerase chain reaction or culture of bacilli from aspirates. But considering the high tubercular disease burden and limited resources, in our population, the presence of epithelioid cell granuloma is considered as an evidence of tuberculous lymphadenitis [6, 7]. With this background, we did a retrospective study with the following aims and objectives: 1) To study the epidemiological pattern of tuberculous lymphadenitis in our population. 2) To study the spectrum of various cytomorphological patterns of Tuberculous lymphadenitis in our population and 3) To correlate the AFB positivity with cytomorphological patterns.


## MATERIALS AND METHODS

This study was carried out in the Department of a tertiary care hospital catering medical services to the rural population. This is a retrospective review of FNAC of lymph node lesions between Oct 2012 and Feb 2015. During this period, FNAC was performed on 891 cases of palpable/ deep lymphadenopathy. Detailed clinical history and meticulous physical examination including the duration of swelling, size, number, matted/ discrete, consistency and mobility were taken into account. Informed consent of patient was taken and in case of minor, consent of parent/ guardian was obtained. FNAC was performed using 22-24G needles attached to 20c.c syringes. One to two passes were given and the aspirated material was smeared onto glass slides. Smears were fixed in 95% ethyl alcohol and stained with Papanicolaou stain. Leishman stain was done on air dried smears. Ziehl-Neelsen(ZN) staining was done whenever a cytological diagnosis of granulomatous disease was made and also in cases with abundant necrosis and suppuration. In cases where fluid was aspirated on FNAC, the fluid was centrifuged and smears were prepared from the sediment followed by the above staining methods. The

cytological diagnosis for each case was based on cytomorphology and available clinical information. For study purpose, the tuberculous lymphadenitis cases were further divided into three groups on the basis of cytomorphological analysis: group 1- Epithelioid cell granulomas with/ without Langhan's giant cells without necrotic material (Fig. 1a), group 2- Epithelioid cell granulomas with/ without Langhan's giant cells with necrotic material (Fig. 1b & 1c), and group 3- Mainly necrotic material with many polymorphs without epithelioid cells or giant cells (Fig. 1d). A definitive cytological diagnosis of TB lymphadenitis could be considered in the smears with the first two patterns, while in the third, with the help of ZN stain (Fig. 2) and relevant routine investigations like Complete blood counts (CBC), Erythrocyte sedimentation rate (ESR) and special investigations like Montoux test, Chest radiograph and Ultrasound of lymphnodes, final diagnosis was made. Data was entered in excel spread sheet and statistical analysis was done. Chi square test was applied to correlate cytomorphological pattern and bacillary load. P value <0.05 was taken to represent significant difference.



**Fig. 1a-** Microphotograph showing epithelioid cell granuloma (Leishman stain; 400x). **Fig. 1b and 1c-** Microphotograph showing epithelioid cell granulomas with necrosis (Leishman stain; 200x). **Fig. 1d-** Microphotograph showing few inflammatory cells in a necrotic background. (Leishman stain; 100x)



**Fig. 2- Acid fast bacilli (Ziehl –Neelsen stain; 1000x)**

**RESULTS**

A total of 891 cases of palpable/deep lymphadenopathy subjected to FNAC were studied. Tuberculous lymphadenitis was the most frequent diagnosis and was seen in 500 cases. Cervical lymph nodes were involved in 418 of 500 cases (83.6%) followed by axillary lymph nodes in 44 cases (8.8%), supraclavicular lymph nodes in 15 cases (3%), inguinal lymph nodes in 12 cases (2.4%). One case (0.2%) of tuberculous mesenteric lymphadenitis was reported (Table 1). Multiple lymph nodes were involved in 10

cases (2%). The age range of Tuberculous lymphadenitis patients was 9 months to 85 years with the male to female ratio of 0.65: 1. Maximum numbers of cases were seen in the age group 10-19 years followed by 20-29 years (Table 2). The size of the lymph nodes varied from 0.8cm to 5cm. The lymph nodes were multiple, soft to firm, matted in 368 cases (73.6%) whereas in 132 cases (26.4%), a single discrete node was seen. Discharging skin sinus was seen in association with matted lymph nodes in 40 cases (8%). Aspirate was blood mixed in 190 cases whereas it was purulent to cheesy in rest of 310 cases. Blood mixed aspirate was predominantly seen in group 1 cases. Family history of tuberculosis was found in 97 cases (19.4%) and 6% cases were defaulters who were diagnosed tuberculosis patients but had not completed ATT course. The TB lymphadenitis cases were further divided into three groups on the basis of cytomorphological analysis; Group 1: Granulomas without necrosis- 133/ 500 (26.6%), Group 2: Caseating epithelioid granuloma- 330/ 500 (66%), Group 3: Necrotizing lymphadenitis- 37/ 500 (7.4%). ZN positivity for acid- fast bacilli was found in 108 cases (21.6%). Different patterns showed varied AFB positivity. Necrotizing lymphadenitis showed 81.0% (30/ 37) positivity. The pattern Caseating epithelioid granuloma had 20.6% (68/330) AFB positivity and in Granuloma without necrosis, it was 7.51% (10/ 133) (Table3). Significant statistical difference was seen among the three patterns and AFB positivity. The p value was <0.001 and was highly significant statistically.

**Table -1: Lymphnode groups involved in Tuberculous Lymphadenitis**

Cervical	418
Axillary	44
Supraclavicular	15
Inguinal	12
Mesenteric	1
Multiple	10

**Table-2: Incidence of Tuberculous lymphadenitis in relation to age and sex**

Age Group (Years)	Tuberculous lymphadenitis	
	Male	Female
0 – 9	33	33
10 – 19	55	103
20 – 29	66	89
30 – 39	21	45
40 – 49	13	19
50 – 59	6	8
60 – 69	4	5
70 – 79	2	0
80 – 89	0	1
	197	303

**Table-3: Various cytomorphological patterns of tuberculous lymphadenopathy**

Cytomorphological picture	No. of cases	AFB positive cases	P value
Granuloma without necrosis	133 (26.6%)	10 (7.51%)	<0.001* Highly significant
Caseating Granuloma	330 (66%)	68 (20.6%)	
Necrotising Lymphadenitis	37 (7.4%)	30 (81.0%)	
Total	500	108	

\*Chi square test was applied.

**DISCUSSION**

FNAC is an important diagnostic tool to aid in the diagnosis of lymph node lesions. It is rapid, inexpensive, safe and reduces the need for surgical biopsy [8]. It was Dudgeon and Patrick in 1927, who first used FNAC in diagnosing Tuberculous lymphadenitis [9]. In our hospital, a rural tertiary care centre, tuberculous lymphadenitis constituted 56.11% of all lymph node aspirates forming the most common cause of lymph node enlargement. Maximum numbers of cases in our study were in the age group 10-19 years followed by the age groups 20-29 years and 30-39 years. Majority of cases in our study (75.2%) were younger than 30 years of age. Whereas majority of patients were in the second to fourth decades of life in studies by Paliwal et al [10], Ergete and Bekele et al [11], Purohit et al [12], Dandapat et al [13], Ahmad et al [14] and Hemlata et al [15]. A declining trend was seen in the geriatric population in concordance with that observed by Ahmed et al [14]. It may be attributed to development of cell mediated immunity against tubercle bacilli in the elderly. Significant female preponderance; 0.65: 1 (M: F) sex ratio was observed in our study. This may be because of malnutrition and overall low living standards among females in this area. Female preponderance has also been observed by Fatima et al [8], Paliwal et al [10], Ergete and Bekele et al [11], Purohit et al [12], Aslam S et al [16], Chand et al [17] and Pamra et al [18]. Male predominance was noted by Ahmad et al [14] and Rajsekaran et al [19]. Cervical lymph nodes (83.6%) were most frequently involved in our study. Similar observation was made by Paliwal et al [10], Pavithra et al [16], Chand et al [17], Khajuria et al [20] and Das et al [21]. In our study, the commonest cause of lymphadenopathy in pediatric age group males was reactive lymphadenitis whereas it was tuberculous lymphadenitis in pediatric age group females. This is in concordance to study by Sharma et al [22] in pediatric age group that showed female predominance and most common involvement of cervical region (88.2%). In our study, matted lymph nodes were seen in 73.6% cases. Ahmed et al [14] also reported that majority of cases (60%) had matted lymph nodes in their study. We noted slightly higher incidence (56.11%) of tuberculous lymphadenitis than that reported by Paliwal et al [10] (55%) from a referral centre for tuberculosis cases. There are wide variations in the prevalence of TB lymphadenitis which may be due to variation in geographical prevalence of TB, age and immunological status of the patients selected. Sarda et al [23] reported

86% of the enlarged nodes to be tuberculous in origin. Similar high incidence was reported by Jha et al [24] (63%), Mahajan et al [25] (54%), Dandapat et al [13] (41.5%), Castro et al [26] (46%), Ahmad et al [14] (38%) and Tilak et al [27] (38.8%). ZN staining for acid fast bacilli was seen in 21.6% cases in our study while Paliwal et al [10] reported 71%, Ergete and Bekele et al [11] reported 71.7%, Chand et al [17] reported 44.54%, Bezabih et al [28] reported 59.5%, Dasgupta et al [29] reported 45.6% AFB positivity in their studies. On the other hand, a low positivity rate of AFB (19.6% cases) on ZN Staining was reported by Aggarwal et al [30] in their study. Low incidence of AFB positivity in our study may be due to combined maximum percentage (92.6%) of cases having epithelioid cell granulomas with necrosis and cases with epithelioid cell granulomas without necrosis unlike series of Paliwal et al [10] where these groups together accounted for only 30.7%. In our study, necrosis alone was seen in 37 cases and 81% of these were positive for acid fast bacilli. It has been observed that the foci of necrosis are associated with marked proliferation of tubercle bacilli whereas lymphocytes, epithelioid cells and multinucleated giant cells have a role in limiting the proliferation of AFB [21, 31]. Therefore, it is expected that FNAC from a tuberculous abscess yields more AFBs than early tuberculous lymphnodes [32]. The characteristic necrotic background comprising of eosinophilic granular material containing nuclear debris was described as 'tubercular diathesis'. Those cases lacking the typical finding and showing scattered epithelioid cells with or without granulomas or only necrotic material with neutrophilic infiltration were diagnosed as tuberculous lymphadenitis when this tubercular diathesis was found cytologically, even though AFB were absent in these smears [33]. Granulomas can be seen in a variety of other conditions causing lymphadenopathy including sarcoidosis, carcinoma, lymphoma, fungal diseases, cat scratch disease, collagen vascular disease and disease of the reticuloendothelial system [34]. But, in a region where tuberculous infection is common and other granulomatous diseases are rare, the presence of a granulomatous feature on FNAC is highly suggestive of tuberculosis [32]. Also for AFB positivity, 10,000-100,000 mycobacterium/ ml of sample should be present for smear AFB positivity [10].

## CONCLUSION

FNAC is a simple Outpatient department (OPD) procedure which is well accepted by patients. It can be effectively used for diagnosing tuberculous lymphadenitis and the diagnostic accuracy can be enhanced by combining cytomorphology with ZN staining. Our study revealed a huge burden of tuberculous lymphadenitis in this region and also highlighted the various cytological patterns of tuberculous lymphadenitis with AFB positivity in these patterns.

## REFERENCES

1. Sharma SK, Mohan A; Extrapulmonary tuberculosis. *Indian J Med Res*, 2004;120:316-53.
2. Shrivastav A, Shah HA, Shrivastav G, Santwani PM; Utility of acid- fast staining and re-aspiration in tubercular lymphadenopathy- 3 year study at tertiary centre. *Int J Med Sci Public Health*, 2013;2:875-9.
3. Pavithra P, Geetha JP; Role of Fine Needle Aspiration Cytology in the evaluation of the spectrum of lymph node lesions. *Int J Pharm Bio Sci*, 2014;5(4):377-84.
4. Corbett EL, Watt CJ, Walker N, Meher D, Williams BG, Raviglione MC et al; The growing burden of tuberculosis: Global trends and interactions with the HIV epidemic. *Arch Intern Med*, 2003;163:1009-21.
5. Laishram RS, Devi RKB, Konjengbam R, Devi RKT, Sharma LD; Aspiration cytology for the diagnosis of tuberculous lymphadenitis: A five year study. *J Indian Acad Clin Med*, 2010;11:31-5.
6. Mohapatra PR, Janmeja AK; Tuberculous lymphadenitis. *J Assoc Physicians India*, 2009;57:585-90.
7. Pandit AA, Khilnani PH, Prayag AS; Tuberculous lymphadenitis: Extended cytomorphologic features. *Diag Cytopathol*, 1995;12:23-7.
8. Fatima S, Arshad S, Ahmed Z, Hasan SH; Spectrum of cytological findings in patients with Neck lymphadenopathy- Experience in a tertiary hospital in Pakistan. *Asian Pacific J Cancer Prev*, 2011;12:1873-5.
9. Singh JP, Chaturvedi NK, Das A; Role of Fine needle aspiration cytology in the diagnosis of tuberculous lymphadenitis. *Indian J Pathol Microbiol*, 1989;32:100-4.
10. Paliwal N, Thakur S, Mullick S, Gupta K; FNAC in Tuberculous lymphadenitis: Experience from a Tertiary Level Referral Centre. *Indian J Tuberc*, 2011;58:102-7.
11. Ergete W and Bekele A; Acid fast bacilli in aspiration smears from tuberculous patients. *Ethiop J Health Dev*, 2000;14(1).
12. Purohit MR, Mustafa T, Morkve O, Sviland L; Gender differences in the clinical diagnosis of tuberculous lymphadenitis- a hospital based study from central India. *International Journal of Infectious Diseases*, 2009;13(5):600-5.
13. Dandapat MC, Panda BK, Patra AK, Acharya N; Diagnosis of tubercular lymphadenitis by fine needle aspiration cytology. *Indian J Tuberc*, 1987;37:139-42.
14. Ahmed SS, Akhtar S, Akhtar K, Naseem S, Mansoor T, Khalil S; Incidence of Tuberculosis from study of Fine- Needle Aspiration Cytology in Lymphadenopathy and Acid- Fast Staining. *Ind J Com Med*, 2005;30(2):63-6.
15. Hemlatha A, Shruti PS, Kumar MU, Bhaskaran A; Cytomorphological patterns of Tubercular Lymphadenitis Revisited. *Ann Med Health Sci Res*, 2014;4(3):393-6.
16. Nataraj G, Kurup S, Pandit A, Mehta P; Correlation of fine needle aspiration cytology, smear and culture in tuberculous lymphadenitis: a prospective study. *Journal of postgraduate Medicine*, 2002; 48(2):113.
17. Chand P, Dogra R, Chauhan N, Gupta R, Khare P; Cytological pattern of Tubercular lymphadenopathy on FNAC: Analysis of 550 consecutive cases. *JCDR*, 2014;8(9):16-19.
18. Pamra SR, Baily GV, Gupta SP; Cervical lymphadenopathies. *Indian J Tuberc*, 1987;34:96-100.
19. Rajshekar S, Gunasekaran M, Bhanumati V; Tuberculous cervical lymphadenitis in HIV positive and negative patients. *Ind Jour Tub*, 2001;48:201-4.
20. Khajuria R, Goswami KC, Singh K, Dubey VK; Pattern of lymphadenopathy on fine needle aspiration cytology in Jammu. *JK Science*, 2006;8(3):157-9.
21. Das DK; Fine Needle Aspiration Cytology in the diagnosis of Tuberculous lesions. *Laboratory Medicine*, 2003;31(11):625-32.
22. Sharma S, Sain R, Khalid UK, Singla N, Sharma PP, Behera D; Clinical profile and treatment outcome of tuberculous lymphadenitis in children using DOTS strategy. *Indian J Tuberc*, 2010;57:4-11.
23. Sarda AK, Bal S, Singh MK, Kapur MM; Fine needle aspiration cytology as a preliminary diagnostic procedure for asymptomatic cervical lymphadenopathy. *J Assoc Physicians India*, 1990;38(3):203-205.
24. Jha BC, Dass A, Nagarkar NM, Gupta R, Singhal S; Cervical tuberculous lymphadenopathy: Changing clinical pattern and concepts in management. *Postgrad Med J*, 2001;77(905):185-187.
25. Mahajan M, Hirachand S, Kafle PK, Bista M, Shrestha S, Toran KC et al.; Incidence of tuberculosis in enlarged neck nodes, our experience. *Kathmandu University Medical Journal*, 2009;7(1):54-58.
26. Castro DJ, Hoover L, Castro DJ, Zuckerbraun L. Cervical Mycobacterial Lymphadenitis. Medical vs surgical management. *Arch Otolaryngol* 1985;111(12):816-9.

27. Tilak V, Dhadel AV, Jain R. Fine needle aspiration cytology of head and neck masses. *Ind J Pathol Microbiol* 2002;45(1):23-30.
28. Bezabih M, Marian DW, Selassie SG. Fine needle aspiration cytology of cervical lymphadenopathy with special reference to tuberculosis. *J Indian Med Assoc.* 1994;92:44-6.
29. DasGupta A, Ghosh RN, Poddar AK et al. Fine needle aspiration cytology of cervical lymphadenopathy with special reference to tuberculosis. *J Indian Med Assoc* 1994;92(2):44-60.
30. Aggarwal P, Wali JP, Singh S, HandaR, Wig N, Biswas A. A clinico bacterial study of Peripheral Tuberculous Lymphadenitis. *J Assoc. Physicians India* 2001;49:808-12.
31. Bhardwaj K, Goyal T, Bhardwaj BL. Fine Needle Aspiration Cytology in lymph node disorders with special reference to tuberculosis. *J of Cytology* 2000;17(3):155-9.
32. Lau SK, Wei WI, Hsu C, Engzell UCG. Efficacy of fine needle aspiration cytology in the diagnosis of tuberculous cervical lymphadenopathy. *J Laryngol Otol* 1990;104(1):24-7.
33. Paul PC, Goswami BK, Chakrabarti S, Giri A, Pramnisk R. Fine needle aspiration cytology of lymphnodes- An institutional study of 1448 cases over a five year period. *J of Cytology* 2004;21:187-90.
34. Finfer M, Perchick A, Barstein DE. Fine needle aspiration biopsy diagnosis of syndrome. Tuberculous lymphadenitis in patients with and without the acquired immune deficiency. *Acta Cytol* 1991;35:325-32.