

## **Research Article**

### **Predictive Role of HuR Protein Expression Level in Colonic Tumor**

**Ibrahim Abdulmajeed Altamemi<sup>1</sup>, Rateb F. Abo Khadher<sup>2</sup>**

<sup>1</sup>Department of Microbiology, College of Medicine, Al-Qadisiya University, Iraq

<sup>2</sup>Al-dewanyia Teaching Hospital (Immunology Unit), Iraq

#### **\*Corresponding author**

Ibrahim Abdulmajeed Altamemi

Email: [Ibrahim.altamemi@gmail.com](mailto:Ibrahim.altamemi@gmail.com)

---

**Abstract:** HuR is an mRNA-binding protein. Intracellular localization of HuR is mainly found within nucleus, but it could be translocate between the nucleus and cytoplasm. In the cytoplasm HuR can increase half-life of certain mRNA target. Since cytoplasmic localization of HuR is essential for its activity, thus, HuR translocation in malignant cells could have prognostic indication. In the present study we aimed to evaluate the significance importance of HuR in the aggressiveness of colorectal adenocarcinoma. To achieve this goal, we have investigated its expression level in adenocarcinoma sample from Iraqi patients, through linking its expression with tumor histopathological variables (stage, grade, grade, and lymph node involvement), by using Immunohistochemical staining method. Study done on 40 colorectal cancer samples and their respective resection margins. Present study demonstrated that, the positive expression rate of integrin HuR in non-tumor colorectal mucosa was significantly lower than that of the colorectal cancer (CRC) tissue ( $P < 0.005$ ). Moreover, when CRC samples breakdown according to histopathological variables, significant differences in expression level of HuR protein when compared with different tumor stage, grade, and LN involvement depending on mean expression  $\pm$ SE value ( $P < 0.05$ ,  $P < 0.05$ , and  $P < 0.05$  respectively). Our results show that high cytoplasmic HuR expression is associated with a poor histologic differentiation, large tumor size, and poor prognosis in colorectal adenocarcinoma.

**Keywords:** Colon cancer, HuR, prognosis, tumor progression, HuR translocation, IHC, histopathology

---

#### **INTRODUCTION**

Aberrant expression of gene products is a general feature of tumorigenesis. Gene expression can be regulated at the level of DNA, RNA or the protein itself. RNA regulation occurs through alterations in translational efficiency and in mRNA stability. mRNA stability depends on cis-elements in the RNA and trans-acting factors. A well-studied mRNA instability element is the AU-rich element (ARE) in the 3' untranslated region (3'UTR). HuR, a member of the Hu/ELAV (embryonic lethal abnormal vision family of RNA-binding proteins, can bind ARE-containing mRNAs through its RNA recognition motif. It is postulated that HuR binds these mRNAs in the nucleus and accompanies them into the cytoplasm (i.e. nucleocytoplasmic translocation), thereby protecting the mRNA from degradation, and affecting the protein expression levels of target genes [1]. It has been hypothesized that HuR contributes to neoplastic growth by regulating expression of genes involved in carcinogenesis that harbor an ARE in the 3'UTR, such as COX-2,  $\beta$ -catenin and vascular endothelial growth factor (VEGF) [2,3,4]. HuR is known to be an important post-transcriptional regulator of VEGF. Although the

native half-life of VEGF messenger RNA (mRNA) is under 1 h, the half-life may increase by 2.5–8-fold when VEGF mRNA is bound by HuR [5,6]. This finding, combined with the observation that HuR expression is augmented in times of hypoxic stress, supports the hypothesis that HuR may be an upstream mediator of tumor angiogenesis [7,8]. Although no HuR mutations have been found to be associated with cancer, links between HuR and malignancies of breast, colon, lung, and ovary have been suggested [9]. Importantly, it has been shown in small series that HuR expression tends to correlate with degree of transformation, that is, levels of HuR are lower in normal mucosa than in adenomas, which in turn have lower levels of HuR expression than carcinomas [9, 10]. We were, therefore, interested in determining whether the expression pattern of HuR in colorectal tumors correlated with histological type or grade. In this set of experiments we attempted to investigate whether high cytoplasmic HuR signal was predictive of poor prognosis, as it has been shown in other cancer models.

**PATIENTS AND METHODS**

**Patients and Sampling**

Forty patients with colorectal adenocarcinoma, who were confirmed histopathologically, were included in this study. Their age were ranged from 20- 80 years. Paraffin embedded blocks of tumor and resection margins were retrieved along with the histopathological report of each patient from histopathological laboratory. For staging of the cancer, astler-coller staging system was adopted in this study [11]. In addition, resection margins were confirmed again to be free of malignancy. Adequate thin paraffin embedded sections (5µm thick) of tumor and resection margins were prepared on positively charged slides for the immunohistochemistry Technique (IHC).

**Immunohistochemical Detection of HuR**

A primary monoclonal antibody against HuR(US biological, USA) reacts with it's antigen. biotinylated secondary antibody then reacts with the primary antibody. This is followed by the attachment of an enzyme-conjugated streptavidin to the biotins on the secondary antibody. The enzyme converts a substrate to a colored reaction product. And the procedude has been done according to manufactural instructions (US biological, USA).There after slides were examined by histopathologist under light microscope (40x).

**RESULTS**

**Histopathological Data**

Forty patients with colorectal adenocarcinoma were investigated. According to the histological differentiation, tumors were broken down in to three

groups, well differentiated (WD, n=10), moderately differentiated (MD, n=20), and poorly differentiated (PD, n=10), moreover, patients were further grouped according to their histopathological criteria, as follow: tumor stage (A, n=9, B, n=10, C, n=11, and D, n=10), lymph node involvement (free, n=19, and involved, n=21).

**Tumor sites versus resection margins**

Based on Immunohistochemical staining method, the present study found that, there were significant difference in the mean expression level of HuR protein between tumor sites and their resection margins (79.45714± 2.29, 32.4± 1.39 ), respectively ( P<0.005),(table 1). A noteworthy here is that high intensity of staining is found in the periphery of necrotic foci of the colorectal cancer, labeling with the HuR antibody distinctly intensified in tumor epithelium adjacent to the necrotic foci. HuR, IHC staining in tumor sample and their resection margin are shown in figure 1.

**Correlation among protein expression of HuR with different histopathological variables**

HuR proteinexpression in colorectal adenocarcinoma was analyzed against the different histopathological features of the tumors based on Spearman's statistical correlation. As shown in Table 2, 3, and 4, current study demonstrated that there were significant differences in expression level of HuR protein when compared with different tumor stage, grade, and LN involvement depending on mean expression ±SE value (P< 0.05, P< 0.05, and P<0.05 respectively).

**Table 1: HuR, protein expression in tumor sites and their resection margins, based on t.test**

Sample	No.	HuR Mean ± SE
Normal	40	32.4± 1.392040868
Tumor	40	79.45714± 2.29370
* (P<0.005).		

**Table 2: Expression level of HuR protein along different stage group**


Stage	No.	Mean ± SE
		HuR
A	9	55.16 ± 1.99
B	10	74.40 ± 1.62
C	11	86.82 ± 1.49
D	10	94.87 ± 0.74
LSD value	----	4.522 *
* (P<0.05).		

**Table 3: Expression level of HuR protein along with different tumor grade**

Grade	No.	Mean ± SE
		HuR
WD	10	62.00 ± 4.47
MD	20	78.79 ± 2.72
PD	10	92.00 ± 2.06
LSD value	---	9.849 *
* (P<0.05).		

**Table 4: Effect of LN involvement on HuR expression level**

LN	No.	Mean ± SE
		HuR
Free	19	67.18 ± 2.69
Involved	21	90.21 ± 1.29
LSD value	----	3.124 *
* (P<0.05).		



**Fig.1:Immunohistochemical staining of HuR in colorectal adenocarcinoma section and their resection margins by DAP (brown color) counterstained with hematoxylin. (A) HuR protein expression in high grade tumor cells which show predominant cytoplasmic staining. (B) HuR protein expression in low grade tumor cells which show nuclear stain. (C and D) Resection margin stained with HuR antibody. Magnification power (40X).**

## DISCUSSION

HuR expression was investigated in normal colorectal mucosa, and patients with colorectal adenocarcinomas. We found increasing cytoplasmic HuR immunoreactivity and nucleocytoplasmic translocation of HuR in tumor tissue compared with their respective resection margins ( $79.45714 \pm 2.29$ ,  $32.4 \pm 1.39$ ;  $P < 0.005$ ) However, when HuR expression was analyzed according different tumor histopathological variables, current study showed that HuR protein expression rate was significantly associated with tumor stage, poor differentiation, and lymphoid node invasion, ( $P < 0.05$ ,  $P < 0.05$  &  $P < 0.05$  respectively), in addition to that we demonstrated a cytoplasmic immunoreactivity or translocation gradually increased subsequently along with tumor grade from well differentiated in to poorly differentiated (figure 1), also we noticed same translocation associated with tumor stage progression, and lymph node metastasis. Whereas, nuclear expression of HuR showed to be limited to tumor free resection margin, and tumor with low grade. Our results are in accord with the previous studies that reported that the HuR protein translocates from the nucleus to the cytoplasm during tumorigenesis[12, 13, 14]. Denkert *et al.* found a gradual increase of cytoplasmic HuR expression in subsequent Dukes stages: 38% of Dukes A carcinomas and 50% of Dukes B carcinomas, 58% of Dukes C carcinomas and 85% of Dukes D carcinomas exhibited cytoplasmic HuR expression.8 Taken together, cytoplasmic HuR expression appears associated with progression to metastasis and poor prognosis[15].

Although the ability of HuR to shuttle from the nucleus to cytoplasm is important for mRNA stabilization, the exact mechanism is not known. Several mechanisms for controlling the cellular location have been studied. Some studies have shown that mitogenic-activated protein kinase-2 increased cytoplasmic translocation of HuR and the stability of ARE-containing mRNAs [16]. The translocation of HuR to the cytoplasm can be induced by stress caused by agents such as UV light, DNA damaging agents or T cell activation, while AMP-activated kinase can inhibit the translocation of HuR to the cytoplasm [17, 18, 19]. There are some other mRNAs, such as the mRNAs for tumor necrosis factor  $\alpha$ , cyclin A, cyclin B1, VEGF, and uPAreceptor that are stabilized by HuR [20, 21, 22]. Based on the known functions of HuR, we believe that HuR might play an important role in cell cycle regulation, apoptosis, angiogenesis, inflammation, and tumor growth. Moreover, HuR can be a potential target for molecular tumor therapy with consideration of these multiple effects.

## CONCLUSION

The cytoplasmic expression of HuR may be a part of a regulatory Pathway (s) that controls the

expression of some mRNA target in colorectal cancer. There are several hundred putative targets for HuR; relatively few of these interactions are well characterized[21].We elected to investigate the interaction between HuR and VEGF. Additional studies with a larger number of specimens are required to determine if HuR might be a potential target for tumor control.

## REFERENCES

1. Brennan CM, Steitz JA; HuR and mRNA stability. *Cell Mol Life Sci*, 2001; 58:266-77.
2. Lopez de Silanes I, Fan J, Yang X, Zonderman AB, Potapova O, Pizer ES, Gorospe M; Role of the RNA-binding protein HuR in colon carcinogenesis. *Oncogene*, 2003; 22:7146-54.
3. Dixon DA, Tolley ND, King PH, Nabors LB, McIntyre TM, Zimmerman GA, Prescott SM; Altered expression of the mRNA stability factor HuR promotes cyclooxygenase-2 expression in colon cancer cells. *J Clin Invest*, 2001; 108:1657-65.
4. Denkert C, Koch I, von Keyserlingk N, Noske A, Niesporek S, Dietel M, Weichert W; Expression of the ELAV-like protein HuR in human colon cancer: association with tumor stage and cyclooxygenase-2. *Mod Pathol*, 2006; 19:1261-9.
5. Levy AP, Levy NS, Goldberg MA; Post-transcriptional regulation of vascular endothelial growth factor by hypoxia. *J Biol Chem.*, 1996; 271(5):2746-53.
6. Stein I, Neeman M, Shweiki D, Itin A, Keshet E; Stabilization of vascular endothelial growth factor mRNA by hypoxia and hypoglycemia and coregulation with other ischemia-induced genes. *Mol Cell Biol.*, 1995; 15(10):5363-8.
7. Galbán S, Kuwano Y, Pullmann R, Martindale JL, Kim HH, Lal A, Gorospe M; RNA-binding proteins HuR and PTB promote the translation of hypoxia-inducible factor 1alpha. *Mol Cell Biol.*, 2008;28(1):93-107.
8. Lejbkowitz F, Goldberg-Cohen I, Levy AP; New horizons for VEGF. Is there a role for nuclear localization? *ActaHistochem.*, 2005; 106(6):405-11.
9. de Silanes IL, Fan J, Yang X, Zonderman AB, Potapova O, Pizer ES, Gorospe M; Role of the RNA-binding protein HuR in colon carcinogenesis. *Oncogene*. 2003; 22(46): 7146-54.
10. Lopez de Silanes I, Lal A, Gorospe M; HuR: post-transcriptional paths to malignancy. *RNA Biol*. 2005; 2(1):11-3.
11. Astler VB, Coller FA; The prognostic significance of direct extension of carcinoma of the colon and rectum. *Ann Surg*, 1954; 139: 846-847.
12. Do SI, Do IG, Kim GY, Lee S, Kim YW, Park YK, et al; Correlation between cyclooxygenase-2

- expression and HuR cytoplasmic translocation of breast cancer. *Korean J Pathol.*, 2008; 42:75-80.
13. Mrena J, Wiksten JP, Thiel A, Kokkola A, Pohjola L, Lundin J, et al; Cyclooxygenase-2 is an independent prognostic factor in gastric cancer and its expression is regulated by the messenger RNA stability factor HuR. *Clin Cancer Res.*, 2005; 11:7362-8.
  14. Denkert C, Weichert W, Pest S, Koch I, Licht D, Köbel M, et al; Overexpression of the embryonic-lethal abnormal vision-like protein HuR in ovarian carcinoma is a prognostic factor and is associated with increased cyclooxygenase 2 expression. *Cancer Res.*, 2004; 64:189-95.
  15. Denkert C, Koch I, von Keyserlingk N, Noske A, Niesporek S, Dietel M, Weichert W; Expression of the ELAV-like protein HuR in human colon cancer: association with tumor stage and cyclooxygenase-2. *Mod Pathol*, 2006; 19:1261-9
  16. Subbaramaiah K, Marmo TP, Dixon DA, Dannenberg AJ; Regulation of cyclooxygenase-2 mRNA stability by taxanes: evidence for involvement of p38, MAPKAPK-2, and HuR. *J Biol Chem.*, 2003; 278:37637-47.
  17. Wang W, Furneaux H, Cheng H, Caldwell MC, Hutter D, Liu Y, et al; HuR regulates p21 mRNA stabilization by UV light. *Mol Cell Biol.*, 2000; 20:760-9.
  18. Atasoy U, Watson J, Patel D, Keene JD; ELAV protein HuA (HuR) can redistribute between nucleus and cytoplasm and is upregulated during serum stimulation and T cell activation. *J Cell Sci.*, 1998; 111:3145-56.
  19. Wang W, Fan J, Yang X, Fürer-Galban S, Lopez de Silanes I, von Kobbe C, et al; AMP activated kinase regulates cytoplasmic HuR. *Mol Cell Biol.*, 2002; 22:3425-36.
  20. Wang W, Caldwell MC, Lin S, Furneaux H, Gorospe M; HuR regulates cyclin A and cyclin B1 mRNA stability during cell proliferation. *EMBO J.*, 2000; 19:2340-50.
  21. Wein G, Rössler M, Klug R, Herget T; The 3'-UTR of the mRNA coding for the major protein kinase C substrate MARCKS contains a novel CU-rich element interacting with the mRNA stabilizing factors HuD and HuR. *Eur J Biochem.*, 2003; 270:350-65.
  22. Tran H, Maurer F, Nagamine Y; Stabilization of urokinase and urokinase receptor mRNAs by HuR is linked to its cytoplasmic accumulation induced by activated mitogen-activated protein kinase-activated protein kinase 2. *Mol Cell Biol.*, 2003; 23:7177-88.