

Comparison of Arnett's Dentoskeletal Soft Tissue Norms of Himachali Male Population with South Indian Male Population Using Digital Tracing Method: A Cephalometric Study

Upadhyay Sachin¹, Sharma Aseem^{1*}, Phor Deepak², Didhra Gaurav³, Mandeep Nain⁴, Yatharth Goel⁵

¹Sr.Lecturer, Department of Orthodontics and Dentofacial Orthopaedics, Himachal Institute of Dental Sciences, Paonta Sahib, Himachal Pradesh, India

²Sr, Lecturer, Department of Orthodontics and Dentofacial Orthopaedics, Yamuna Dental College, Yamunanagar, Haryana, India

³M.D.S. (Department of Periodontics and Implantology), Director at Dent-O-Care, Basdehra, Mehatpur, Himachal Pradesh 174315, India

⁴P.G. Student (Department of Prosthodontics), DAV Dental College, Yamunanagar, Haryana, India

⁵Director at Dantam Dental Solutions, Ann's School, 44 Civil Lines, Jadugar Road, Near St, Roorkee, Uttarakhand 247667, India

DOI: 10.36347/sjds.2019.v06i09.011

| Received: 13.09.2019 | Accepted: 21.09.2019 | Published: 30.09.2019

*Corresponding author: Dr. Aseem Sharma

Abstract

Original Research Article

Background: This analysis was developed directly from the philosophy expressed in Arnett and Bergman "Facial keys to orthodontic diagnosis and treatment planning, Parts I and II". This article describes comparison of various dentoskeletal traits between Himachali male population and South Indian male population that contribute to an aesthetically pleasing face which should be considered during orthodontic treatment. The aim of the present study was to highlight the differences in dentoskeletal norms between Himachali males and South Indian males. **Materials and Methods:** Lateral cephalograms of 50 Himachali males of age group 18-25 years were taken in Department of Orthodontics, Himachal Institute of Dental Sciences, Paonta Sahib and tracing of soft tissue profile as well as related osseous and dental structures were made using Nemoceph software. Then Arnett's dentoskeletal traits were compared with Himachali traits. **Results:** The present study showed that there is increased overjet in Himachali population but the soft tissue thickness of upper lip, lower lip and pogonion is more in South Indian population. **Conclusion:** Individual norms are necessary for a population in order to plan and deliver quality treatment.

Keywords: Himachali males, South Indian males, dentoskeletal traits, treatment planning.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Preservation of facial attractiveness is an important goal of orthodontic treatment. Due to a complicated interaction of genetic and environmental factors the morphological features of an individual vary from race to race. Even within the same race, each subgroup had its own standards. Hence, the established norms for other ethnic group can not apply to the population of Himachal Pradesh. Therefore, the applicability of various dentoskeletal parameters proposed by Bergman, should be analyzed which will improve treatment planning for male population of Himachal Pradesh [1]. As Soft tissue norms serve as a guideline in calculating change it has been suggested that certain cephalometric standards relating teeth to cranial or facial bones could ensure good facial form if adhered to as a treatment goal [2]. The attainment of facial soft-tissue proportionality is one of the principal goals in the treatment of dentofacial deformities and can be achieved with properly planned and executed orthognathic surgery techniques [3]. Therefore the aim

of this study is to compare the dentoskeletal norms for the male population of Himachal Pradesh with the South Indian male population so that the soft tissue cephalometric values of various parameters could be differentiated so as to guide the orthodontist towards a better diagnosis and treatment planning of dentofacial deformities for the population of Himachal Pradesh.

MATERIALS AND METHODS

This study was carried out in the Department of Orthodontics and Dentofacial Orthopaedics of Himachal Institute of Dental Sciences, Paonta Sahib (H.P). 50 Males in the age group of 18-25 years who were residents of Himachal Pradesh were considered for the study. All subjects were examined by a panel of the faculty members of the Department of Orthodontics and Dentofacial Orthopaedics and reasonably balanced faces were selected. Small diameters silver beads of dimension 2 mm (approx.) were chosen for the use as metallic markers.

All the cephalograms of patients were taken in natural head position with Carestream X-ray machine with model no CS8100 (2016). All the radiographs were traced using (Nemoceph) Dental Studio NX (2006) software after the calibration of the images.

All the landmarks were marked manually using inbuilt autozoom feature of the software and the measurement of the values and was done automatically by the Nemoceph software. Ten radiographs were also retraced after one week to check the intra-operator

reliability. The Arnett's soft tissue cephalometric analysis (STCA) [4] was performed considering:

Dentoskeletal Factors

- Maxillary occlusal plane (Mx OP) to true vertical line (TVL)
- Maxillary central incisor tip (Mx1) to maxillary occlusal plane (Mx OP)
- Mandibular incisor tip (Md1) to mandibular occlusal plane (Ms OP)
- Overjet (OJ)
- Overbite (OB)




Fig-1: Dentoskeletal factors

Values of all parameters thus obtained were organized in the form of a master chart using Microsoft Excel and were statistically analyzed and compared with South Indian male population.

OBSERVATIONS AND RESULTS

Comparison of Himachali males with South Indian value males shows that there are significant

differences in values for Himachali ethnic male population and South Indian male population.

In the dentoskeletal factors the value for overjet was 3.83 ± 0.91 mm for Himachali males and 2.90 ± 0.88 mm for South Indian males and the difference in mean was significant with p value of <0.001 (Table-1).

Table-1: Comparison of Cephalometric Parameters between Himachali and South Indian Males

	Himachali Males		South Indian Value Males		t Value	p Value
	Mean	S.D.	Mean	S.D.		
Mx1-MxOP	55.14	5.305	54.47	5.71	0.877	0.379
Overjet	3.836	.9194	2.90	0.88	7.199	<0.001**
Md1-Md OP	62.878	7.0245	64.27	-15.60	-1.401	0.167
Overbite	2.782	1.6834	3.18	0.96	-1.672	.101

Statistically significant value at *p ≤ 0.05, **p ≤ 0.01 and ***p ≤ 0.001

DISCUSSION

The perception of beauty is different for different people. There are no rules governing why a face is beautiful. As professionals have increased their ability to change faces, the necessity to understand what

is and is not beautiful has increased [5]. Angle suggested that if teeth were placed in optimal occlusion, good facial harmony would result [2]. The facial skeleton and its overlying soft tissue determine facial harmony and balance. It is the structure of the overlying

soft tissues and their relative proportions that provide the visual impact of the face [1].

The primary goal of treatment becomes soft tissue harmony and balance, not Angle's ideal occlusion. Keeping this in mind while planning treatment is very important [6].

The Soft Tissue Cephalometric Analysis is a radiographic instrument that represents the clinical extension of the philosophy detailed in "Facial keys to orthodontic diagnosis and treatment planning." This cephalometric soft tissue analysis guides soft tissue examination, as do these earlier articles, but with added advantages. Because the STCA is a cephalometric analysis, profile soft tissue landmarks are easily seen, marked, and measured cephalometrically. Importantly, the midface metallic markers, for the first time, allow important soft tissues (orbital rim, cheekbone, subpupil, and alar base) to be easily seen, marked, and measured [4].

The STCA has five distinct but cross-contributory elements:

- First, the system analyzes key dentoskeletal structures controlled by the orthodontist (Mx1 to MxOP, Md1 to MdOP) and surgeon (MxOP to TVL). Orthodontic and surgical manipulation of the dentoskeletal factors is key to facial profile and esthetics.
- Second, it measures key soft tissue structures that affect facial appearance.
- Third, it measures important vertical soft tissue lengths and soft tissue to hard tissue relationships.
- Fourth, it measures soft tissue points relative to the TVL, thus producing absolute projection values for each point.
- Fifth, the absolute values are then related to one another to test facial harmony. Harmony numbers provide a test of facial balance within the individual's face and, importantly, are independent of the true vertical anteroposterior placement.

The present study was carried out in the Department of Orthodontics and Dentofacial Orthopaedics of Himachal Institute of Dental Sciences, Paonta Sahib (H.P). A sample of 50 Males who were residents of Himachal Pradesh in the age group of 18-25 years were considered for the study. The subjects were first assessed clinically, in natural head position, seated condyles, and with lips at rest. Then, facial examination (frontal/profile) was used as described by Arnett and Bergman [7] with particular emphasis on midface structures that do not show on standard cephalometric analysis. In particular, orbital rim, subpupil and alar base contours were noted to indicate anteroposterior position of the maxilla.

Small diameters silver beads of dimension 2 mm (approx.) were chosen for the use as metallic markers on the basis of their excellent radio – opaque properties even in small dimensions and since they are light weight they could easily applied on the face to the precise location with the help of paper tape according to Arnett *et al.*, [4] in 1999. Next in the preparation for cephalometric radiograph, metallic markers in the form of small beads of silver were placed on the right side of the face to mark key mid face structures.

With the midface structures marked, the Natural Head Position was recorded. The subjects were asked to swallow and bite into centric occlusion. A cephalogram was obtained with subjects positioned in natural head position, seated condyle, and with lips at rest. The natural head position was recorded based on the method proposed by Cooke and Wei in the year 1988 according to which the subject tilted the head forward and backward with decreasing amplitude until a comfortable position of natural balance was achieved.

With the mirror, the subject was then requested to look into the reflection of their eyes in a mirror located 200 cm ahead. Special care was taken to ensure that the head was not moved when the ear posts were carefully inserted.

The digital cephalograms obtained were then traced with the help of (Nemotec) Dental Studio 2006 software. After the cephalometric soft and hard tissue landmarks were measured on 50 facially balanced males of Himachali origin, dentoskeletal, soft tissue, vertical, projection and facial harmony norms and SDs were established. The values were obtained and statistical analysis was done with Student's t test.

The STCA integrates occlusal correction and soft tissue balance. Of all the STCA [4] measurements, only five relate hard tissue to hard tissue points: maxillary occlusal plane, maxillary incisor to maxillary occlusal plane, mandibular incisor to mandibular occlusal plane, overbite, and overjet. These hard tissue relationships are measured because to a large extent they control the esthetic outcome of occlusal treatment [4].

The STCA is not meant as a stand-alone cephalometric analysis. It is meant to be used in combination with clinical facial examination and cephalometric treatment planning, to provide clinically relevant soft tissue information with checks and balances (between cephalometric and clinical facial findings) [4].

This study highlights the differences in dentoskeletal structures of Himachali and South Indian male population [8]. Himachali males and South Indian males show significant differences in certain parameters

like there is increased overjet in Himachali males as compared to South Indian males.

These findings show that group specific norms are an essential prerequisite for accurate evaluation of orthodontic patients. The attainment of facial soft-tissue proportionality is one of the principal goals in the treatment of dentofacial deformities and can be achieved with properly planned and executed orthognathic surgery techniques. It would also help to evaluate the progress of treatment in various stages.

CONCLUSION

All male patients cannot be treated to one set of norms as the facial skeleton and its overlying soft tissue is different for Himachali and South Indian population. So, it is of utmost importance to maintain the facial harmony and balance of a particular population while treating a patient of that particular population.

REFERENCES

1. Burstone CJ. Lip posture and its significance in treatment planning. American Journal of Orthodontics. 1967 Apr 1;53(4):262-284.
2. Spyropoulos MN, Halazonetis DJ. Significance of the soft tissue profile on facial esthetics. American Journal of Orthodontics and Dentofacial Orthopedics. 2001 May 1;119(5):464-71.
3. Proffit WR, Fields Jr HW, Sarver DM. Sarver. Contemporary Orthodontics. 2013; 4.
4. Arnett GW, Jelic JS, Kim J, Cummings DR, Beress A, Worley Jr CM, Chung B, Bergman R. Soft tissue cephalometric analysis: diagnosis and treatment planning of dentofacial deformity. American Journal of Orthodontics and Dentofacial Orthopedics. 1999 Sep 1;116(3):239-53.
5. Scheideman GB, Bell WH, Legan HL, Finn RA, Reisch JS. Cephalometric analysis of dentofacial normals. American journal of orthodontics. 1980 Oct 1;78(4):404-420.
6. Burstone CJ. The Integumental profile. Am Journal of Orthodontics. 1958; 44(1):1-25.
7. Arnett GW, Bergman RT. Facial keys to orthodontic diagnosis and treatment planning. Part I. American journal of orthodontics and dentofacial orthopedics. 1993 Apr 1;103(4):299-312.
8. Kalha AS, Latif A, Govardhan SN. Soft-tissue cephalometric norms in a South Indian ethnic population. American journal of orthodontics and dentofacial orthopedics. 2008 Jun 1;133(6):876-81.