

Original Research Article

A study of biochemical changes in gallstone disease among alcoholics in karaikal population

Maheshwari Narayanan^{*1}, R.Bharathidasan², Manoji³, K.Kalaiselvi⁴, Thangapaneerselvam⁵, G.Ambujam⁶

¹Post graduate, ³Assistant professor, ⁴Professor and HOD, ⁵Professor, Department of biochemistry, Vinayaka Mission Medical College and Hospital, Karaikal-609609, Puducherry, India

²Assistant professor, ⁶Professor and HOD, Department of surgery, Vinayaka Mission Medical College and Hospital, Karaikal-609609, Puducherry, India

*Corresponding author

Maheshwari Narayanan

Email: maheshwari22.narayanan@gmail.com

Abstract: Gallstones are the major cause of morbidity worldwide, Gallstones (cholelithiasis) develops because of an imbalance in the chemical composition of bile inside the gallbladder and it is due to multifactorial etiology which includes chronic alcoholism. Patients undergoing (cholecystectomy) once symptoms appear. The important complications of long term gallstones are acute pancreatitis, cholecystitis, obstructive jaundice which alter the liver function and decrease the life expectancy. The main aim in the present study was conducted to assess the Biochemical variations in gallstone diseases among alcoholics in coastal region of karaikal. The Materials and methods were A total of 22 patients(case) in which 11 alcoholics and 11 non alcoholics who were age and sex matched with multiple stones in gall bladder, symptomatic and undergoing cholecystectomy were recruited for this study . 25 normal individuals as controls were recruited for this study. Serum total and direct bilirubin were estimated by DMSO method. Serum SGPT, SGOT, GGT, Lipase, Amylase and Serum alkaline phosphates level were estimated. The Results In the present study liver function biochemical parameters were found to increase in alcoholic gallstone disease individuals when compared to healthy individuals and also compared to gallstone disease nonalcoholic subjects. ALP and Direct bilirubin were significantly elevated in alcoholic gallstone disease subjects ($p=0.01$, $p=0.04$) when compared to healthy individuals. Finally the Conclusion thus elevated biochemical parameter levels in blood could predict the presence of liver damage and also indicates the severity of gallstone disease and its complications among alcoholics in coastal region.

Keywords: Gallstones (cholelithiasis), SGPT, SGOT, GGT, Lipase

INTRODUCTION:

Gallstones are among the most common gastrointestinal disorders in Western populations. Approximately 80 percent of gallstones contain cholesterol (as cholesterol monohydrate crystals). The remaining 20 percent are pigment stones, which consist mainly of calcium bilirubinate and will not be discussed in this article [1]. Cholesterol-containing gallstones are divided into two subtypes: cholesterol stones (which contain 90- to 100-percent cholesterol) and mixed stones (which contain 50- to 90-percent cholesterol). Each sub type may also contain varying amounts of calcium salts, bile acids, and other components of bile. Cholelithiasis (gallstone formation) results from a combination of several factors, including super saturation of bile with cholesterol, accelerated nucleation of cholesterol monohydrate in bile, and bile stasis or delayed gallbladder emptying due to impaired

gallbladder motility [2]. Cholesterol super saturation can result from an excessive concentration of cholesterol in bile, a deficiency of substances that keep cholesterol in solution (i.e., bile salts and phospholipids), or a combination of these factors. While most gallstones are asymptomatic, some patients experience biliary colic, which is characterized by sudden and severe right-upper-quadrant pain (often accompanied by nausea and vomiting), occurring postprandially and lasting one to four hours. Acute or chronic cholecystitis may also occur in association with gallstones. Complications of cholecystitis may include infection, perforation, and gangrene [3]. Cholecystectomy is the most frequently recommended conventional treatment for symptomatic gallstones. Bile acids (ursodeoxycholic acid or chenodeoxycholic acid) are also used in some cases to dissolve radiolucent stones, but these drugs can cause gastrointestinal side

effects and there is a high rate of stone recurrence after treatment is discontinued. Lithotripsy is used in some cases in conjunction with ursodeoxycholic acid for patients who have a single symptomatic non-calcified gallstone. There is evidence that dietary factors influence the risk of developing cholesterol gallstones [4].

MATERIALS AND METHODS:

A total of 22 patients (case) in which 11alcoholics and 11 non alcoholics who were age and sex matched with multiple stones in gall bladder, symptomatic and undergoing cholecystectomy were recruited for this study. 25 normal individuals as

controls were recruited for this study. Serum total and direct bilirubin were estimated by DMSO method. Serum SGPT, SGOT, GGT, Lipase, Amylase and Serum alkaline phosphates level were estimated for research purpose in-patients gastroenterological department of vinayaka mission university karikal. To verify the diagnosis of diabetes and NASH, determining the functional state of the liver using complex clinical laboratory, biochemical and instrumental methods. Statistical analysis Results were expressed as mean ± SD and mean ± SEM. Students t test was used to compare the data between patients and control groups.

RESULTS:

Table 1: show the comparison of plasma biochemical changes between case or group A and control group B

Parameter	Case Patient with gallstone disease	Control Healthy individuals	P value
Total bilirubin (mg/dl)	0.93	0.52	< 0.001*
Direct bilirubin (mg/dl)	0.62	0.32	< 0.001*
SGPT IU/L)	38.4	33	< 0.001*
SGOT (IU/L)	34.1	32.3	< 0.001*
ALP IU/L)	153	110	<0.001*
GGT (IU/L)	75.4	35.2	0.002*
LIPASE (IU/L)	61.9	42.1	< 0.01*
AMYLASE (IU/L)	94.2	75.2	< 0.01*

Legend : 1 All the parameter are increased in group A (case) when compared to group B(control)and direct bilirubin ,alkaline phosphatase(ALP) and gamma glutamyl transepeptidase(GGT) were significantly increases in group B when compared to group A

Table 2: show the comparison of plasma biochemical changes between group 1(Alcoholic) and group (2) (non alcoholic) Gallstone diseased individuals in group A

Parameter	Group 1	Group 2	P value
Total bilirubin (mg/dl)	0.93	0.42	0.006
Direct bilirubin (mg/dl)	0.61	0.3	0.002*
SGPT (IU/L)	36.2	33.1	0.003*
SGOT (IU/L)	35.6	32.1	0.006
ALP (IU/L)	163	98	<0.001*
GGT (IU/L)	98	76.3	0.002*
LIPASE (IU/L)	59.3	32.1	0.006
AMYLASE(IU/L)	93.2	73.2	0.006

Legend: 2 All the parameter are increased in group1 when compared to group 2 , and direct bilirubin ,alkaline phosphatase(ALP) , SGPT and gammaglutamyltrance peptidase(GGT) were significantly increases in group 1(alcoholic) then group 2 (non alcoholic)

DISCUSSION

We discovered that gallstone disease affects the course of nonalcoholic steatohepatitis and accompanied by decompensation of carbohydrate, protein and lipid metabolism, as in patients of the third study group (nonalcoholic steatohepatitis without pathology of the gallbladder) changes in clinical and laboratory parameters were the most insignificant compared to first group and second group [5]. The amount of cholesterol in the bile is supposed to increase

with age. This is caused by dyslipoproteinemia that results in a linear increase in cholesterol excretion into the bile and by the reduced synthesis of bile acids due to the dropped activity of the enzyme cholesterol 7 α -hydroxylase (CYP7A1) [6]. The xenobiotic receptor, pregnant X receptor (PXR), has a role in the pathogenesis of cholesterol GD. PXR prevents cholesterol GD *via* its coordinated regulation of the biosynthesis and transport of bile salts in the liver and intestine. Cholesterol precipitation is prevented by

increases in concentrations of biliary bile salts and a reduced cholesterol saturation index (CSI). Loss of PXR sensitized mice to lithogenic diet-induced cholesterol GD, characterized by decreases in biliary concentrations of bile salts and phospholipids and increases in the CSI and formation of cholesterol crystals [7]. The decreased bile acid pool size in PXR-/- mice that received lithogenic diets was associated with reduced expression of CYP7A1, the rate-limiting enzyme of cholesterol catabolism and bile acid formation. The reduced expression of CYP7A1 most likely resulted from activation of PXR and induction of fibroblast growth factor 15 in the intestine. Normal bile consists of 70% bile salts (mainly cholic and chenodeoxycholic acids), 22% phospholipids (lecithin), 4% cholesterol, 3% proteins, and 0.3% bilirubin. An increased expression of gel-forming mucin, such as MUC5AC and MUC2, was found in patients with hepatolithiasis [8] Giannini E and coworkers described a positive correlation between MUC1 and MUC5AC expression, indicating a gene-gene interaction that might affect the accumulation of mucin gel and cholesterol GS formation. Bile mucin is derived from pure hepatic bile, gallbladder-concentrated bile, and mucin secreted by the bile duct epithelium. In patients with biliary sludge, mucin concentration was higher in bile collected by endoscopic retrograde cholangiography than in gallbladder bile. Cholestasis enhances the synthesis and release of ALP, and accumulating bile salts increase its release from the cell surface. ALP half-life in the circulation is about 1 week [4]. These characteristics explain why ALP levels usually rise late in bile duct obstruction and decrease slowly after resolution [9]. Elevated GGT levels can be observed in a variety of hepatic diseases, including chronic obstructive pulmonary disease and renal failure, and may be present for weeks after acute myocardial infarction. Increased serum levels observed in alcoholic liver disease can be the result of enzyme induction and decreased clearance [10].

CONCLUSION:

Alterations in liver enzyme levels are one of the most common problems encountered in everyday clinical practice. Finding the way through the multiple diagnostic pathways can challenge even the experienced clinician. The pattern of enzyme abnormality, interpreted in the context of the patient's characteristics, can aid in directing the subsequent diagnostic work-up.

REFERENCES:

1. Bosma PJ, Chowdury JR, Bakker C, Gantla S, de Boer A, Oostra BA, et al.; The genetic bases of the reduced expression of bilirubin UDP-glucuronosyltransferase 1 in Gilbert's syndrome. *N Engl J Med*, 1995; 333(18):1171-5.
2. Thomsen HF, Hardt F, Juhl E; Diagnosis of Gilbert's syndrome. *Scand J Gastroenterol*, 1981; 16(5):699-703.
3. Van Hootehem P, Fevery J, Blanckaert N; Serum bilirubins in hepatobiliary disease: comparison with other liver function tests and changes in the postobstructive period. *Hepatology*, 1985; 5(1):112-7.
4. Doumas BT, Peters T; Serum and urine albumin: a progress report on their measurement and clinical significance. *Clin Chim Acta*, 1997; 258(1):3-20.
5. Pugh RN, Murray-Lyon IM, Dawson JL, Pietroni MC, Williams R; Transection of the oesophagus for bleeding oesophageal varices. *Br J Surg*, 1973; 60 (8): 646-9. [
6. Bonis PA, Tong MJ, Blatt LM, Conrad A, Griffith JL; A predictive model for the development of hepatocellular carcinoma, liver failure, or liver transplantation for patients presenting to clinic with chronic hepatitis C. *Am J Gastroenterol*, 1999;94(6):1605-12.
7. Kamath PS, Wiesner RH, Malinchoc M, Kremers W, Therneau TM, Kosberg CL, et al.; A model to predict survival in patients with end-stage liver disease. *Hepatology*, 2001; 33(2):464-70.
8. Williams AL, Hoofnagle JH; Ratio of serum aspartate to alanine aminotransferase in chronic hepatitis. Relationship to cirrhosis. *Gastroenterology*, 1988; 95 (3): 734-9
9. Giannini E, Botta F, Fasoli A, Ceppa P, Risso D, Lantieri PB, et al.; Progressive liver functional impairment is associated with an increase in AST/ALT ratio. *Dig Dis Sci*, 1999; 44(6):1249-53.
10. Giannini E, Risso D, Botta F, Chiarbonello B, Fasoli A, Malfatti F, et al.; Validity and clinical utility of the aspartate aminotransferase-alanine aminotransferase ratio in assessing disease severity and prognosis in patients with hepatitis C virus-related chronic liver disease. *Arch Intern Med*, 2003; 163(2):218-24.