

Assessment of Cardiovascular Risk Factors in Rheumatoid Arthritis Patients Using Framingham Risk Score

Chakraborty R^{1*}, Haque MS¹, Islam MT¹, Rahman MR², Yesmin S³, Mazumder SK⁴, Sarker RSC⁵

¹Registrar, Department of Internal Medicine, BIRDEM General Hospital, Dhaka, Bangladesh

²Professor & Head, Department of Internal Medicine, BIRDEM General Hospital, Dhaka, Bangladesh

³Associate Professor & Head, Department of Rheumatology, BIRDEM General Hospital, Dhaka, Bangladesh

⁴Junior Consultant, Department of Rheumatology, BIRDEM General Hospital, Dhaka, Bangladesh

⁵Assistant Professor, Department of Internal Medicine, BIRDEM General Hospital, Dhaka, Bangladesh

DOI: <https://doi.org/10.36347/sasjm.2025.v1i04.020>

| Received: 10.03.2025 | Accepted: 16.04.2025 | Published: 22.04.2025

*Corresponding author: Ranodhir Chakraborty

Registrar, Department of Internal Medicine, BIRDEM General Hospital, Dhaka, Bangladesh. Email: cranodhir@gmail.com

Abstract

Original Research Article

Background: Chronic autoimmune rheumatoid arthritis generates harm to body systems beyond joints since it increases patients' vulnerability to developing cardiovascular disease. There are no studied reports examining how to evaluate cardiovascular risks for RA patients in Bangladesh. **Objectives:** The purpose of this study is to determine the cardiovascular risk factors among RA patients by applying the Framingham Risk Score (FRS) to identify principal risk predictors. **Methodology:** A two-year cross-sectional study took place at BIRDEM General Hospital between July 2022 and June 2024. A risk score analysis according to the Framingham model required data from eight criteria, including the patient's age, together with cholesterol measurements and blood pressure readings and smoking status, and presence of diabetes. According to Framingham Risk Score classification, patients were divided into low risk (<10%) and intermediate risk (10–20%), and high risk (>20%). SPSS version 25.0 was used for statistical analysis under a $p < 0.05$ significance level. **Result:** The research included 100 RA patients with 78% females and 22% males, who fulfilled specified enrollment requirements. Analysis revealed that FRS assessment categories divided the study participants as 41% low risk while 36% had intermediate risk and 23% displayed high risk levels. This study established diabetes mellitus ($p < 0.001$), dyslipidemia ($p = 0.001$), hypertension ($p = 0.003$), increasing age ($p < 0.001$), and elevated C-reactive protein (CRP) levels ($p = 0.002$) as major risk factors that increased FRS. The participants with RA that exceeded 10 years of existence demonstrated an increased cardiovascular risk profile, but this correlation did not meet statistical significance ($p = 0.009$). Research findings supported earlier studies which showed that RA affects females at a ratio of 3.5 to 1 than males. **Conclusion:** The study demonstrates that patients with RA exhibit a substantial rise in their CVD risk, and diabetes, dyslipidemia, and hypertension serve as major causal elements. These findings underline the need for RA patients to receive regular cardiovascular screening as well as comprehensive preventive measures, including lifestyle improvements and proper healthcare management protocols.

Keywords: Cardiovascular risk, Rheumatoid Arthritis Framingham Risk Score.

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Rheumatoid Arthritis (RA) is a chronic systemic autoimmune disease that initially causes joint inflammation, pain, swelling, and stiffness but damages other tissues and organs, including the heart, kidneys, lungs, digestive system, eyes, skin, and nervous system, after a certain period [1-3]. According to the WHO, 18 million people are suffering from RA worldwide as of 2019 [4]. With widespread inflammation, RA generates systemic complications [5]. Researchers suggested that rheumatoid arthritis increased the risk of cardiovascular

disease 1.5-2 folds [6]. Hypertension, diabetes, obesity, smoking, and an unhealthy lifestyle increase cardiovascular risks, along with rheumatoid arthritis [7]. The concept of Framingham Risk Score (FRS) was first developed in 1948 by Framingham Heart Study by a long-term epidemiological study [8]. It is a calculation tool that predicts an individual's risk of developing cardiovascular disease by analyzing related risk factors like age, sex, cholesterol levels, blood pressure, smoking status, and diabetes [9]. According to previous research, 4.5% of Bangladeshi patients suffer from cardiovascular disease [10]. The prevalence of rheumatoid arthritis

Citation: Chakraborty R, Haque MS, Islam MT, Rahman MR, Yesmin S, Mazumder SK, Sarker RSC. Assessment of Cardiovascular Risk Factors in Rheumatoid Arthritis Patients Using Framingham Risk Score. SAS J Med, 2025 Apr 11(4): 368-372.

among cardiovascular patients is unexplored in Bangladeshi settings. This study aims to determine the cardiovascular risk factors among rheumatoid arthritis patients by using the Framingham Risk Score.

METHODOLOGY

A bi-year cross-sectional study was conducted in the Department of Internal Medicine, BIRDEM General Hospital, in the timeline of July 2022 to June 2024. Framingham Risk Score was calculated based on the Age (years), Total Cholesterol (mg/dL), HDL Cholesterol (mg/dL), Systolic Blood Pressure (SBP) (mmHg), Hypertension Treatment (Yes/No), Smoking Status (Current/Former/Never), Diabetes Status (Yes/No) of patients and the interpretation was given as Low Risk (<10%), Intermediate Risk (10–20%) and High Risk (>20%) [11]. Patients' data were collected by following the declaration of Helsinki and some study-specific inclusion and exclusion criteria.

Inclusion criteria

- Patient age ≥ 18 years.

- Confirmed diagnosis of Rheumatoid arthritis by blood tests: Rheumatoid Factor (RF) and Anti-CCP Antibody.
- Willingness to provide written informed consent.

Exclusion criteria

- Pre-existing cardiovascular disorder.
- Known or self-claimed pregnancy.
- Chronic kidney disease.
- Severe hepatic impairment.

The researcher collected data by using individual case report forms (CRFs) and after a successful data collection, all data were analyzed by MS Excel and SPSS version 25.0. A p -value<0.05 was considered significant in a 95% confidence interval.

RESULT

The study enrolled a sum of 100 patients, including 78 females and 22 males. The female-to-male ratio calculated for this study is 3.5:1.




Figure 1: Pie chart showed demographic distribution of patients, sex (N=100)

Figure 01 shows the prevalence of female participants over male participants in this study. The

female to male ratio proves that females are at three times more risk than male patients.




Figure 2: Column chart showed demographic distribution of patients, age (N=100)

Figure 02 portrays that the majority of the patients diagnosed with rheumatoid arthritis are between the ages of 40-49 years. The mean age determined by

analysis was 45.8 ± 11.6 years, with an upper age limit of 71 years and lower limit of 33 years.

Table I: Clinical Characteristics of patients with rheumatoid arthritis (N=100)

Variable	RA Patients (n=100)
Hypertension, n (%)	42 (42.0%)
Diabetes Mellitus, n (%)	20 (20.0%)
Dyslipidemia, n (%)	48 (48.0%)
Smoking Status, n (%)	
- Current Smoker	15 (15.0%)
- Former Smoker	18 (18.0%)
- Never Smoked	67 (67.0%)
BMI (kg/m ²), mean \pm SD	27.3 ± 4.0
Family History of CVD, n (%)	28 (28.0%)

Table I showcases the clinical profile of study subjects that describes that 42% of patients have existing hypertension, 20% diabetes mellitus, and 48% Dyslipidemia. The percentage of former and current

smoker patients is 33%. The mean BMI calculated 27.3 ± 4.0 kg/m², which indicates overweight characteristics, and a total 28% of study subjects were found with a family history of cardiovascular diseases.

Table II: Distribution of patients by Framingham Risk Score (N=100)

Risk Category	Number of Patients (n)	Percentage (%)
Low Risk (<10%)	41	41.0%
Intermediate Risk (10–20%)	36	36.0%
High Risk (>20%)	23	23.0%

Table II categorizes patients by Framingham Risk Score, which indicated that 23% of patients are at

high risk of CVD, 41% are at low risk, and 36% are at intermediate risk.

Table III: Key predictors of increase FRS among RA patients (N=100)

Variable	FRS Increase (%)	p-value
Diabetes Mellitus (DM)	+9.2	<0.001*
Dyslipidemia	+7.5	0.001*
Hypertension	+6.3	0.003*
RA >10 years	+5.1	0.009
Age (per decade)	+4.8	<0.001*
Elevated CRP	+3.9	0.002*

Table III draws a relationship between clinical factors and FRS among rheumatoid arthritis patients. Data shows that diabetes mellitus significantly increased FRS ($P < 0.001$). Dyslipidemia, Hypertension, Aging, and elevated CRP also contributed to increased risk of CVD ($P < 0.05$). Rheumatoid arthritis duration of more than 10 years is not clinically significant to the association of increase Framingham Risk Score.

DISCUSSION

Framingham Risk Score (FRS) uses a specific algorithm to calculate and predict cardiovascular risk factors in patients. In North America, this calculation is measured as a validated predictor for CVD risk [12]. In Bangladesh, previous studies have been carried out to predict CVD risk in diabetic patients [13]. This study assessed cardiovascular risk factors among rheumatoid arthritis patients by using Framingham Risk Scoring. A combined 100 subjects with RA were included in this study, where the female-male ratio was 3.5:1. A study on

rheumatoid arthritis shows the female/male ratio to be 3:1 in rheumatoid arthritis which is similar to recent findings [14]. Kvien T K *et al.*, 2006 published that females are more prone to rheumatoid arthritis than males, and he also added the rate incidence is 4-5 times higher below the age of 50 [15]. The current study resulted in the mean age of patients being 45.8 ± 11.6 with a range of 33-71 years, where the majority of the cases were detected in the age of 40-49 years. An Egyptian cohort study on rheumatoid arthritis calculated the mean age for male subjects to be 44.9 ± 10.2 years and for female patients 49.9 ± 13.2 years; 83.1% of their study subjects were female, and 16.2% were male [16]. This study and the previous studies suggest that women after 30 years of age are at more risk of rheumatoid arthritis. Due to increase hormonal fluctuation [17], especially estrogen influence [18], genetic (X-linked) factors [19], and microchimerism [20], RA is more common in females than males. The clinical profile of RA patients reveals that 42% of patients are suffering

from hypertension, 20% diabetes mellitus, and 48% dyslipidemia. Earlier research demonstrated that hypertension is 32.35% common in patients with rheumatoid arthritis [21]; DM incidents rates in patients with RA is 5.2% to 16.7% according to studies [22]; Prevalence of dyslipidemia is 56.5% in rheumatoid arthritis patient [23]. Dyslipidemia refers to a condition of abnormal rising of lipid levels in blood [24]. Guimarães M F B R *et al.*, 2019 disclose the tends to grow obesity in RA patients and the mean BMI calculated by their study was $27.1 \pm 4.9 \text{ kg/m}^2$ [25] which is similar findings to this study ($27.3 \pm 4.0 \text{ kg/m}^2$). Considering related factors, this study predicted that 41% of patients are at low risk of cardiovascular disease, 36% of patients are at intermediate risk, and 23% of patients are at high risk of CVD. Islam A M *et al.*, 2016 found that the rate of rheumatoid heart disease and stroke in Bangladesh is 20-25% [26], which is agreeable to the study findings. The $p\text{-value} < 0.001$ suggests that DM in RA patients exerts a potential risk to develop cardiovascular disorder. Previous research also suggested DM tends to obesity, insulin resistance, and uncontrolled hyperglycemia, which increase the risks of cardiovascular disease [27]. More preventive care for patients with DM and RA should be taken to lower the risk of CVD [28]. Dyslipidemia and Hypertension are some other significant factors ($p < 0.05$) to increase Framingham Risk Score to develop CVD by the study findings. Lifestyle modification and opting for a healthy diet with low sodium is advisable to control dyslipidemia and hypertension in RA patients [29]. The risk of CVD significantly rises ($p < 0.05$) with increasing age, and elevated CRP indicates inflammation in body, which is also related to developing CVD ($p < 0.05$) in rheumatoid arthritis patients. Regular exercise, weight loss, and dietary control can decrease inflammation, as well as C-reactive protein, and lower the risk of cardiovascular disease [30].

LIMITATION

The single-center study with a limited population does not generalize the overall rheumatoid arthritis patients in Bangladesh. A details research with long-term follow-up will provide the fact-based outcome of Framingham Risk Score to predict the development of cardiovascular disease in rheumatoid arthritis patients at Bangladeshi settings.

CONCLUSION

The research demonstrates that higher cardiovascular risk exists for RA patients due to the presence of diabetes along with dyslipidemia and hypertension. The Framingham Risk Score effectively categorized cardiovascular disease risk for this study group. Regular screening in combination with early intervention through proper medical care as well as lifestyle changes, must become essential practices to help RA patients with their significant cardiovascular risks.

Author Contribution

Conceptualization, Methodology, Formal analysis, Investigation, Data collection, analysis, and original draft by RC.

MSH, MTI, MRR, SY, SKM, and RSCS participated in writing, reviewing and editing, and visualizing.

Conceptualization, Methodology, Formal analysis, Investigation, Data collection, analysis, and original draft by Chakraborty R.

Haque MS, Islam MT, Rahman MR, Yesmin S, Mazumder SK, Sarker RSC participated in writing, reviewing and editing, and visualizing.

Abbreviations

RA: Rheumatoid Arthritis

FRS: Framingham Risk Score

CVD: Cardiovascular Disease

DM: Diabetes Mellitus

HDL: High-density lipoprotein

SBP: Systolic Blood Pressure

CRP: C-reactive Protein

BMI: Body Mass Index

CRFs: Case Report Forms

WHO: World Health Organization

Conflicts of Interest: The authors declare no conflicts of interest.

REFERENCES

1. Krati Chauhan, J. S.-D. (2023). Rheumatoid Arthritis. In Treasure Island (FL). StatPearls Publishing. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK441999/>
2. V. F. A. M. Derksen, T. W. (2017). The role of autoantibodies in the pathophysiology of rheumatoid arthritis. *Semin Immunopathol*, 39, 437–446. doi: <https://doi.org/10.1007/s00281-017-0627-z>
3. Giles, J. T. (2019). Extra-articular Manifestations and Comorbidity in Rheumatoid Arthritis: Potential Impact of Pre-Rheumatoid Arthritis Prevention. *Clinical Therapeutics*, 41(7), 1246 - 1255. doi:10.1016/j.clinthera.2019.04.018
4. WHO. (2023). Rheumatoid arthritis. World Health Organization. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/rheumatoid-arthritis>
5. Di Wu, Y. L. (2022). Systemic complications of rheumatoid arthritis: Focus on pathogenesis and treatment. *Front Immunol.*, 13, 105108. doi:10.3389/fimmu.2022.1051082
6. Holly John, G. K. (2009). Cardiovascular comorbidity in early rheumatoid arthritis. *Best Pract Res Clin Rheumatol.*, 23(1), 71-82. doi: 10.1016/j.berh.2008.11.007
7. Nishant Johri, S. V. (2023). Association of cardiovascular risks in rheumatoid arthritis patients: Management, treatment and future perspectives. *Health Sciences Review*, 8, 100108. doi:

- <https://doi.org/10.1016/j.hsr.2023.100108>
8. Rie Sasaki-Nakashima, T. K. (2017). Successful prediction of cardiovascular risk by new non-invasive vascular indexes using suprasystolic cuff oscillometric waveform analysis. *Journal of Cardiology*, 69(1), 30-37. doi: <https://doi.org/10.1016/j.jjcc.2016.06.004>
 9. C Sohn, J. K. (2012). The framingham risk score, diet, and inflammatory markers in Korean men with metabolic syndrome. *Nutr Res Pract.*, 6(3), 246–53.
 10. Fouzia Khanam, M. B. (2019). Prevalence and Risk Factors of Cardiovascular Diseases among Bangladeshi Adults: Findings from a Cross-sectional Study. *J Epidemiol Glob Health.*, 9(3), 176-184. doi:10.2991/jegh.k.190531.001
 11. Mostafa Abohelwa, J. K. (2023). The Framingham Study on Cardiovascular Disease Risk and Stress-Defenses: A Historical Review. *J. Vasc. Dis.*, 2(1), 122-164. doi: <https://doi.org/10.3390/jvd2010010>
 12. Bosomworth, N. J. (2011). Practical use of the Framingham risk score in primary prevention. *Can Fam Physician.*, 57(4), 417–423.
 13. Akter, N. (2020). Prediction of 10-Year Risk of Cardiovascular Disease among Diabetic Patients in Bangladesh. *Delta Medical College Journal*, 8(1), 32 – 40. doi: <https://doi.org/10.3329/dmcj.v8i1.58961>
 14. Beatrice Maranini, A. B. (2022). Focus on Sex and Gender: What We Need to Know in the Management of Rheumatoid Arthritis. *J. Pers. Med.*, 12(3), 499. doi: <https://doi.org/10.3390/jpm12030499>
 15. Tore K Kvien, T. U. (2006, Jun). Epidemiological aspects of rheumatoid arthritis: the sex ratio. *Ann N Y Acad Sci.*, 1069, 212-22. doi:10.1196/annals.1351.019
 16. NESREEN S. ABD EL-GHANY, I. M. (2019). Gender Impact on Rheumatoid Arthritis Disease Characteristics in A Cohort of Egyptian Patients. *Med. J. Cairo Univ.*, 87(3), 1895-1899. Retrieved from <https://www.medicaljournalofcairouniversity.net/>
 17. M Gerosa, V. D. (2008). Rheumatoid arthritis: a female challenge. *Womens Health (Lond).*, 4(2), 195-201. doi:10.2217/17455057.4.2.195
 18. Maria Fernanda Romo-García, M. Z.-Z.-M.-D. (2020). The Role of Estrogens in Rheumatoid Arthritis Physiopathology. In *Rheumatoid Arthritis - Other Perspectives towards a Better Practice*. doi:10.5772/intechopen.81489
 19. Vollenhoven, R. F. (2009). Sex differences in rheumatoid arthritis: more than meets the eye... *BMC Med.*, 7(12). doi:10.1186/1741-7015-7-12
 20. Jacqueline E Oliver, A. J. (2009). Why are women predisposed to autoimmune rheumatic diseases? *Arthritis Res Ther.*, 11(5), 252. doi:10.1186/ar2825
 21. Al-Ahmari, A. K. (2022). Prevalence of Hypertension and Its Associated Risk Factors Among Patients with Rheumatoid Arthritis in the Kingdom of Saudi Arabia. *International Journal of General Medicine*, 15, 6507—6517. doi: <https://doi.org/10.2147/IJGM.S370956>
 22. Jad Okais, F. F. (2022). Association between Diabetes and Rheumatoid Arthritis: A Systematic Literature Review. *The Open Rheumatology Journal*, 16. doi:10.2174/18743129-v16-e221205-2022-13
 23. Mio Akiyama, T. M.-i. (2015). Prevalence of dyslipidemia in Japanese patients with rheumatoid arthritis and effects of atorvastatin treatment. *Clin Rheumatol.*, 34(11), 1867-75. doi:10.1007/s10067-015-3049-0
 24. Jiahui Yan, S. Y. (2023). Dyslipidemia in rheumatoid arthritis: the possible mechanisms. *Front Immunol.*, 14, 1254753. doi:10.3389/fimmu.2023.1254753
 25. Maria Fernanda Brandão de Resende Guimarães, C. E. (2019). High prevalence of obesity in rheumatoid arthritis patients: association with disease activity, hypertension, dyslipidemia and diabetes, a multi-center study. *Advances in Rheumatology*, 59(44). doi: <https://doi.org/10.1186/s42358-019-0089-1>
 26. AKM Monwarul Islam, A. M. (2016). Cardiovascular Disease in Bangladesh: A Review. *Bangladesh Heart Journal*, 31(2). doi: <https://doi.org/10.3329/bhj.v31i2.32379>
 27. Fabio Cacciapaglia, F. R. (2023). Clinical Features of Diabetes Mellitus on Rheumatoid Arthritis: Data from the Cardiovascular Obesity and Rheumatic DISease (CORDIS) Study Group. *J Clin Med.*, 12(6), 2148. doi:10.3390/jcm12062148
 28. E. Ikdahl, S. R. (2020). THU0123 DIABETES MELLITUS AND CARDIOVASCULAR RISK MANAGEMENT IN PATIENTS WITH RHEUMATOID ARTHRITIS IN A LARGE INTERNATIONAL AUDIT. *Annals of the Rheumatic Diseases*, 79(1), 276-277. doi: <https://doi.org/10.1136/annrheumdis-2020-eular.5398>
 29. Panniyammakal Jeemon, R. G. (2017). Management of Hypertension and Dyslipidemia for Primary Prevention of Cardiovascular Disease. In *Cardiovascular, Respiratory, and Related Disorders (3rd ed.)*. Washington (DC): The International Bank for Reconstruction and Development/ The World Bank. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK525163/#>
 30. Ridker, P. M. (2005). C-Reactive Protein, Inflammation, and Cardiovascular Disease. *Tex Heart Inst J.*, 32(3), 384–386.